

Effects of supplementing mid-lactation dairy cows with seaweed and vitamin E on plasma and milk α -tocopherol and antibody response to immunization

Short title: *seaweed and vitamin E supplements to dairy cows*

A. KIDANE¹, I. L. NESHEIM^{2†}, H. J. S. LARSEN³, E. THUEN², S. K. JENSEN⁴ AND H. STEINSHAMN^{1*}

¹ *Bioforsk, Norwegian Institute for Agricultural and Environmental Research, Organic Food and Farming Division, Gunnars veg 6, N-6630 Tingvoll, Norway*

² *Department of Animal and Aquacultural Sciences, Norwegian University of Life Sciences, P.O. Box 5025, N-1432 Ås, Norway*

³ *Department of Food Safety and Infection Biology, Norwegian University of Life Sciences, P.O. Box 8156, Dep., N-0033 Oslo, Norway*

⁴ *Department of Animal Science, Aarhus University, DK-8830 Tjele, Denmark*

(MS received 12 September 2014, revised 5 December 2014, accepted TBC January 2015)

SUMMARY

The objective of the current experiment was to compare the effects of supplementing mid-lactation dairy cows with *all-rac*- α -tocopheryl acetate (*SyntvE*), *RRR*- α -tocopheryl acetate (*NatvE*) or seaweed meal (*seaweed*) in the presence of a *control* group (no supplemental vitamin E or seaweed) on the concentration of α -tocopherol in plasma and milk, and antibody response following immunization. The hypothesis was that supplementation of dairy cows with vitamin E, regardless

*To whom all correspondence should be addressed. Email: havard.steinshamn@bioforsk.no. Telephone: +47 404 80 314.

†Present address: Norsk Landbruksrådgiving Rogaland, Postvegen 211, Særheim, 4353 Klepp st., Norway

of its form, would increase plasma and milk α -tocopherol compared to the control diet and this incremental response would be bigger with *NatvE* than *SyntvE*. Furthermore, it was hypothesized that vitamin E, regardless of its form, will provide an improved adaptive immune response to immunization than the control diet, and cows supplemented with seaweed meal would produce better adaptive immune response following immunization than cows in the control group. Twenty four Norwegian Red (NR) dairy cows in their mid-lactation were allocated randomly to the four treatments in a replicated Latin square design. The cows were fed on a basal diet of silage and concentrate on top of which the experimental supplements were provided. Plasma and milk α -tocopherol concentrations were higher in *NatvE* and *SyntvE* groups than in the other two groups. The RRR- α -tocopherol stereoisomer was the predominant form (> 0.86), in both plasma and milk, whereas the remaining part was largely made up of the other three 2R stereoisomers (RRS, RSR and RSS). In cows fed the *Control*, *seaweed* and *NatvE*, the proportion of the RRR- α -tocopherol stereoisomer in plasma and milk constituted < 0.97 of the total α -tocopherol. Mid-lactation NR dairy cows had higher than adequate levels of plasma α -tocopherol (9.99 mg/l) even when not supplemented with external source of vitamin E, suggesting that with a good quality silage these cows may not be at risk of vitamin E deficiency. Furthermore, the study shows that dairy cows in mid to late lactation have preferential uptake of RRR stereoisomer of α -tocopherol compared to other stereoisomers. All cows responded well to immunization with different antigens, but there were no significant group effects of the diet on the immune response measured.

INTRODUCTION

Self-sufficiency and recirculation of nutrients within the farm are central elements in the organic principles (European Commission 1991; IFOAM 2007). As such, necessary supplements such as vitamins for farm animals under organic settings should be of natural sources if possible (European

Commission 1999). When deemed to be insufficient, extra vitamin E is supplemented in animal feeds as an antioxidant and also because of its diverse physiological roles (Dersjant-Li & Peisker 2010). Some of the functions of vitamin E are effects against testicular degeneration, muscle dystrophy, resorption of foetus and lipid peroxidation (Bouwstra *et al.* 2008; Jensen & Lauridsen 2007). Vitamin E is also important for ruminant meat and milk quality to protect membrane lipids and myoglobin from oxidation (Liu *et al.* 1995; Lee *et al.* 2003; Al Mabruk *et al.* 2004).

Dairy cow diets contain different levels of vitamin E depending on type of feed and the proportion of ingredients in the diet (Beeckman *et al.* 2010; NRC 2001). Therefore, supplemental sources are mandatory when the levels are below what is required by animals for a given production. The additives exist mainly in two commercially available forms of vitamin E; namely, RRR- α -tocopheryl acetate (natural, derived from vegetable oil) and *all-rac- α* -tocopheryl acetate (synthetic, α -tocopherol produced by chemical synthesis) (Dersjant-Li & Peisker 2010). It is well established that RRR- α -tocopheryl acetate has higher bioavailability than *all-rac- α* -tocopheryl acetate in peri-parturient dairy cows (Meglia *et al.* 2006; Weiss *et al.* 2009), but less is known about the availability of the tocopherol-isomers in mid and late lactating cows. Still, the recommended intake of total (supplemental plus feedstuff origin) vitamin E, i.e. 2.6 IU/kg of body weight (NRC 2001), is the same for cows during late gestation and in lactation.

Recent surveys and experiments have indicated that the diets of organic winter-fed dairy cows have varying α -tocopherol contents and that cows, particularly during peripartum, have low plasma vitamin E concentration (Govasmark *et al.* 2005; Sivertsen *et al.* 2005; Beeckman *et al.* 2010; Lindqvist *et al.* 2011; Mogensen *et al.* 2012). As such, it is recommended that diets of dairy cows on organic farms should be supplemented with vitamin E. However, in organic farming, all necessary supplementation to the basic diets should ideally be of natural origin (European Commission 1999). As there are few or no alternatives, the Nordic countries have a present

derogation from the EU regulations and may use synthetic vitamin A, D and E (European Commission 2005). Thus, it is deemed important to look for natural anti-oxidant sources to replace the synthetic ones.

Recent work has suggested that supplementation with a meal prepared from the macro algae *Ascophyllum nodosum*, a seaweed, or treatment of endophyte-infested pasture with *A. nodosum* product had positive effects on product quality (improved beef shelf-life) and stress tolerance (improved immune function) (Allen *et al.* 2001). Alginates containing high proportions of mannuronic acid polymers, such as *A. nodosum*, have been documented to be non-specific immunostimulants, through stimulation of the cytokine production by human monocytes (Otterlei *et al.* 1991) and by increasing the superoxide production in macrophages from fish *in vitro* (Rokstad *et al.* 1996) and *in vivo* (Skjermo & Bergh 2004). Historically, *A. nodosum* has been used as supplementary feed for ruminants in Norway and is currently harvested, dried and milled commercially and sold as a feed supplement. A dairy cow experiment in the late 1960s demonstrated improved performance (+ 6 % in total milk yield) when lactating dairy cows were supplemented daily with 160 g *A. nodosum* (Jensen *et al.* 1968). However, the effects of *A. nodosum* supplementation in mid-lactation dairy cows on the adaptive immune response has so far remained elusive.

The objective of the current study was to test the consequences of feeding two different sources (forms) of supplemental vitamin E (RRR- α -tocopheryl acetate = *NatvE* or all-rac- α -tocopheryl acetate = *SyntvE*) on α -tocopherol and its stereoisomer content of plasma and milk and also on immune response of mid-lactation dairy cows relative to seaweed meal and control. The hypotheses were that supplementation of dairy cows with vitamin E, regardless of its form, would increase plasma and milk α -tocopherol compared to the non-supplemented cows and this increment would be bigger with *NatvE* than *SyntvE*. Furthermore, supplementation with vitamin E, regardless

of its form, would provide an improved adaptive immune response to immunization compared to the control group, and cows given the seaweed meal supplementation would produce better adaptive immune response following immunization than cows in the control group.

MATERIALS AND METHODS

Animals, treatments and experimental design

The experiment was conducted from 22 August 2011 to 9 December 2011 with 24 (16 primiparous and 8 multiparous) Norwegian Red cows (NR) in their mid-lactation period at the start of the experiment (164 d in milk, SD = 30; BW=511 kg, SD=64 kg) at the Animal Production Experimental Centre, Norwegian University of Life Sciences, facility at Aas, Norway. The experiment was carried out in a 4×4 Latin square design (four groups over four periods) with six replicates (squares). During each of the experimental period that lasted for 4 weeks, the first 3 weeks were used for adaptation and the last week was used for sampling. Cows within each of the six squares were matched for age, stage of lactation and level of production.

The experiment was carried out according to the laws and regulations controlling experiments on live animals in Norway, the Norwegian Animal Research Authority, Oslo, Norway.

Feeds and chemical composition of feeds

Cows grazed a summer pasture before the commencement of the experiment and were provided with supplements of silage and concentrates produced for the experiment in the last week of grazing as an adaptation. Mineral and vitamin supplements were also given during the adaptation week.

Basal feed

During the experimental period the cows were fed on silage and concentrates that were produced according to standards for organic production (European Commission 1991). The silage was prepared from the first and second cut of a grass/clover ley preserved in round bales and fed *ad libitum* as an equal mixture of the two cuts through individual feeding gates, and the concentrate was given at a flat rate of about 3.0 kg/cow per day through computerized dispensers. Silage samples were taken for analysis each day of the sampling week by hand collection of representative amount immediately after offering.

Experimental supplements

In addition to the basal feed, the cows in each group were fed one of the following four experimental supplements: *NatvE* (vitamin E added as RRR- α -tocopheryl acetate, Vitfoss A/S, DK-6300 Gråsten, Denmark), *SyntvE* (vitamin E added as *all-rac*- α -tocopheryl acetate, Zhejiang 138 Medicine Co Ltd, Zhejiang, China), *seaweed* (commercially available dried and ground macro algae *A. nodosum*, AlgeaFeed 3.5, Algea AS, Lødingen, Norway) and *control* (with no added vitamin E or macro algae). The *control*, *NatvE* and *syntE* consisted (g/g DM) of barley (0.74), molasses (0.04) and a premix of minerals and vitamin A and D (0.21) supplied by Normin AS (Hønefoss, Norway) (Table 1). The seaweed was a mixture of macro algae (0.30), barley (0.52), molasses (0.04) and minerals (0.14) to achieve recommended supplementations of some important minerals in order to be similar to the other treatments with respect to total mineral supplementation. The *NatvE* and *SyntvE* supplements were formulated to provide 3800 mg α -tocopheryl acetate/kg supplement and an extra supply of 2275 mg α -tocopherol per cow daily. The *control*, *NatvE* and *SyntvE* supplements were offered at a daily rate of 0.7 kg/cow and the *seaweed* supplement at a rate of 0.8 kg/cow in order maintain iso-energetic levels, assuming that the *A. nodosum* meal had a net energy content of 3.9 MJ/kg DM (Jensen *et al.* 1968).

The amount of seaweed meal per cow per day was approximately the same amount as was given in a previous experiment by Jensen *et al.* (1968), where Black Sided Trønderfe and Nordlandsfe (STN) cows were given *c.* 160 g seaweed per day. This corresponded to *c.* 0.35 g seaweed meal per kg body weight. An NR cow weighed between 550 and 650 kg, and therefore was given *c.* 200 g of seaweed meal daily to provide the same amount in relation to body weight as in the experiment by Jensen *et al.* (1968).

The experimental supplements were fed at milking hours, mixed manually with concentrates in the parlour to avoid discriminating refusal by cows. However, any feed refusal (concentrate + experimental supplements) was collected and weighed. The cows had free access to drinking water.

Feed samples were collected daily during the sampling week and mixed to one sample for each feed and period. The samples were stored at -20°C , freeze dried (Christ LCM-2, Beta 1-16 and Christ LOC-1m, Alpha 1-4, Martin Christ, Osterode am Harz, Germany; Hetosicc, Birkerød, Denmark), milled (1.0 mm screen) (Retsch SM 100, Retsch GmbH, Haan, Germany) and stored at -20°C in plastic bags prior to analysis of chemical constituents. The samples were then analysed at Eurofins Food and Agro Testing, Moss, Norway, for dry matter (DM) (European Commission 2009), ash (European Commission 2009), crude protein (CP) (European Commission 2009), and neutral detergent fibre (NDF) was determined with an ANKOM220 fibre analyser (ANKOM Technology, Fairport, NY, USA) according to (Mertens 2002) using sodium sulphite, alpha amylase and ash correction (aNDFom). Undried silage samples (10 g) were homogenized and diluted with 40 ml of deionized water and stored at 4°C for 24 h before pH was measured with a Thermo Orion 420A+ pH-meter with Orion 9107BN electrode (Thermo Scientific, Beverly, MA, USA). Ammonia nitrogen was analysed with MAN-TECH PC-titrate (Guelph, ON, Canada) using an Orion ion analyser 901. Silage samples (20 g) were homogenized and diluted with 40 ml

deionized water and stored frozen, then thawed and filtered before analysing organic acids and ethanol by HPLC using a VA 300/7.8 Nucleogel Ion 300 OA column (Machery-Nagel) at 50°C (mobile phase, 0.010 M H₂SO₄ at 0.6 ml/min) with an UV spectrophotometric detector for lactic acid and a refractive index detector for other acids and ethanol.

For analysis of fatty acids (FA), fat was extracted in a mixture of chloroform and methanol according to Bligh & Dyer (1959) after acidification by boiling in 3 mol/l hydrochloric acid (HCl) for 60 min (Jensen 2008). The FA was analysed as FA methyl esters as described by Jensen & Nielsen (1996) by gas chromatography (Hewlett Packard 6890 series, Agilent Technologies, Palo Alto, CA) equipped with an automatic on-column injector (Hewlett Packard 7673) (Split ratio 4.325:1); a capillary column of 30 m x 320 µm inner diameter; 0.25 µm film thickness (Omegawax, Supelco 4-293-415, Sigma-Aldrich, St. Louis, MO), and a flame ionization detector using C17:0 as an external standard.

For α-tocopherol analysis, 2 g dried material was suspended in 70 ml ethanol, 30 ml methanol, 30 ml ascorbic acid in water (200 g/l) and 20 ml potassium hydroxide (KOH)-water (1:1 w/v). This mixture was saponified for 30 min at 80 °C in the dark and cooled in cold water. Exactly 2 ml of the saponified mixture was diluted in 1 ml water after which tocopherol was extracted twice using 5 ml heptane and analysed by HPLC (Perkin Elmer HS-5-Silica column, 4.0 x 125 mm, Waltham, MA) as described by Jensen & Nielsen (1996) and Jensen *et al.* (1998). Stereoisomers of α-tocopherol were analysed by HPLC as follows: The remaining heptane extract was evaporated to constant dry matter under a stream of nitrogen at 50 °C and 50 µl ethylene glycol dimethyl ether and 25 µl KOK were added to the dried samples. Thereafter the tubes were filled with argon and 45 µl dimethyl sulphate was added. The samples were shaken for 1 h (IKA-VIBRAX-VXR, Janke & Kenkel, Germany), evaporated and 100 µl water and 1.5 ml heptane added. The samples were then vortexed and centrifuged for 10 min at 3000 rpm. The supernatant were transferred to cap

tubes and the stereoisomers determined according to Jensen & Lauridsen (2007).

Response to immunization

All cows were injected subcutaneously with 1mg each of human serum albumin (HSA; Sigma-Aldrich, lot no: 035K7566), ovalbumin (OVA; Sigma-Aldrich, lot no: 120K1201), equine serum albumin (EQSA; BIEWORLD, lot no:ESA62-887) and canine serum albumin (CSA; BIEWORLD, lot no:CASA62-1076) mixed each with Aluminum Hydroxide Gel Adjuvant (Brenntag Biosector, Alhydrogel, lot no: 4601) (1:4) in saline, 6 days before the start of the feeding experiment. Blood was collected from a vein at the tail into five EDTA-tubes from each cow, just before immunization and on day 24 of the first period. Two samples were spun on site and the supernatant plasma was taken and stored frozen at -20°C for later analysis of α -tocopherol. Whole blood for serum collection was sampled at the same time. The serum samples were stored at -20°C until assayed.

The production of antigen-specific antibodies to HAS, OVA, EQSA and CSA was assayed in blood samples in the beginning of the experiment and 24 days following immunization using an enzyme-linked immunosorbent assay (ELISA). Antigen (5 $\mu\text{g/ml}$) diluted in 0.05 M carbonate buffer, pH 9.6, was used for coating of polystyrene microtiter plates (Immunoplates II, Nalge Nunc International, Denmark) by adding 0.2 ml antigen to each well. The plates were sealed, incubated for 3 days at 4°C and washed five times (Microwash II, Skatron, Norway) with phosphate buffered saline (PBS) containing 0.05% Tween 20 (PBS-Tween). One percent skimmed milk (powder, Merck, Germany) in PBS-Tween, 0.2 ml/well, was used to prevent non-specific binding, and the plates were washed five times with PBS-Tween. All test samples were diluted at least 1:100 and further diluted 2-fold in PBS-Tween containing 1% bovine serum albumin (BSA), and 100 μl of

the samples were placed in each well. The plates were sealed and incubated overnight at 4 °C. After incubation, the plates were washed again five times with PBS-Tween, and 100 µl of a 1:5000 dilution (in PBS-Tween with 1% rabbit serum) of a horseradish peroxidase-linked rabbit anti bovine IgG (Sigma, Lot no: 070M4857) was added to each well. The plates were then incubated for 1 h at 37 °C, and washed five times with PBS-Tween. Then 100 µl of the enzyme substrate, ABTS (2,2'-azinobis (3-ethylbenzthiazolinesulfonic acid; Sigma-Aldrich, Lot no: 050M1502V)) was added to each well. The reaction was stopped after 45 min at 37 °C by adding 20 µl sodium azide (NaN₃, 1 mg/ml) and measured immediately with a Titertek Multiskan Plus ELISA reader (Labsystems, Finland; wavelength 405 nm). The final titer was defined as the reciprocal (log₂) of the highest dilution of serum showing positive reaction, i.e. at least 0.1 OD₄₀₅ which exceeded by more than a factor of two of the average OD₄₀₅ of the mean normal control samples in the same test plate.

Milk yield, milk and plasma sampling and chemical analysis

Cows were milked twice daily in a milking parlour, and the amount of milk was recorded automatically at each milking.

Aliquot milk samples were taken during the sampling period from morning and evening milk from each cow four times a week (Monday and Tuesday on each of the two milking times: morning and night). Milk samples intended for analysis of α-tocopherol concentration and FA composition were frozen (−20 °C) immediately. Milk samples intended for analysis of fat and protein were preserved with 2-bromo-2-nitropropane-1,3-diol (Bronopol, D&F Inc., Dublin, CA) and analysed by a Fourier Transformed Infrared Spectroscopy milk analyser (MilkoScan 6000 FTIR, Foss, Hillerød, Denmark).

Before extraction of α -tocopherol, milk and plasma samples were thawed, and milk samples were heated at 40 °C for 20 min and mixed thoroughly prior to analysis. The extraction procedure was as follows: 1 ml of milk, or 500 μ l of plasma, was diluted with 1.0 ml ascorbic acid solution (200 g/l), 0.5 ml methanol, 2.0 ml ethanol, and 0.3 ml KOH-water (1:1 w/v). Saponification was carried out at 80 °C for 20 min. After cooling, tocopherol was extracted twice using 5 ml heptane per occasion. Stereoisomers of α -tocopherol in plasma and milk were determined as described for feed.

Statistical analysis

The data for feed intake, milk and plasma content of α -tocopherol were analysed using the PROC MIXED procedure of SAS (SAS for windows, version 9.2, SAS ltd.) with treatment, period and their interaction as fixed effects and square and cow within a square as random effects. The optimal covariance structure was modelled using an analysis of repeated measures specifying that measures taken on the same cow in subsequent periods are correlated. Satterthwaite's method was used for computing the denominator degrees of freedom.

Antibody titers to the immunization agents were transformed to reciprocal \log_2 units before statistical analysis. Analyses of variance of the response to immunization was performed on specific antibody levels (absorbance) and antibody titers using dietary treatment as fixed effect, the pre-immunization value as covariate and replicate and cow within a replicate as random effects.

Multiple mean comparison was made using Tukey Test at a significance level of 5%. Whilst full *P*-values are presented in the body of the text, only *P*-values below 0.1 are presented in the tables, for clarity and judgment by the reader.

RESULTS

Feed chemical composition and dry matter intake

Chemical composition of the four experimental supplements and the basal feeds are presented in Table 2. Basal feeds (concentrate and silage) were similar for all treatment groups. The chemical composition of the experimental supplements was similar with respect to CP and fat content, but the seaweed supplement had 68 % higher aNDFom content than the other supplements. The concentration of α -tocopherol was 1052 and 993 mg/kg DM in the *NatvE* and *SyntvE* supplements, respectively, which was about 60–70 % less than originally planned, probably due to high temperature during the pelleting process of the supplements (Weiss, unpublished as reported in (NRC 2001)). The content of α -tocopherol in the *control* supplement was higher than expected and it contained also synthetic isomers, which was likely due to contamination from the production of feed previous to the production of the control (Dag Henning Edvardsen, Normin AS, pers. comm.). The silage was well fermented with low concentrations of volatile FA and low proportion of ammonia, and contained 39 mg α -tocopherol/kg DM.

Dry matter intake (DMI, kg/d) is presented in Table 3. There was no difference in total daily DMI among treatment groups (mean \pm SE: 18.1 \pm 0.47). However, mean intake of concentrates did differ ($P = 0.021$) among the treatments due to refusals. Intake of experimental supplements differed among treatments according to design. As such, cows in the *seaweed* group consumed more of the concentrates compared to the *SyntvE* group.

Milk yield and milk fat

Average daily milk production among the treatment groups was not different from each other (Table 4). The control group had higher ($P < 0.05$) mean milk fat content (42.7 g/kg) than either

the *SyntvE* (41.1 g/kg) or the seaweed group (41.2 g/kg). Milk fat content from the *NatvE* treatment was intermediate and not significantly different from the other treatments.

Alpha-tocopherol intake and its stereoisomers in feed, plasma and milk

Total mean daily α -tocopherol intake (g/d) was highest in the *NatvE* group (1146), intermediate in the *SyntvE* (1070) and lowest in the *Control* (636) and *seaweed* (591) groups (Table 3). The latter two groups were not significantly different from each other. As expected, α -tocopherol intake from silage was similar among treatments whilst it was different among treatments based on concentrates source ($P = 0.004$) and based on the experimental supplements ($P < 0.001$) (Table 3).

At the stereoisomer level, the RRR intake was highest in the *NatvE* group, with intermediate and similar intake in the *SyntvE* and *Control* group, whilst the *seaweed* group had the lowest calculated intake. Intake of the rest of the 2R stereoisomers (RRS, RSR and RSS) was also different ($P < 0.001$) among treatments: as such, dairy cows in the *SyntvE* group had the highest intake whilst cows in the *NatvE* group had intermediate and cows in the control and seaweed groups had the lowest intake (Table 3). Similarly, the 2S stereoisomer intake was highest ($P < 0.001$) in the *SyntvE* group being *c.* 0.23 of total α -tocopherol intake. The cows in the *NatvE* group had very low level intake of 2S stereoisomer (~ 0.028 of total α -tocopherol intake) compared to the *SyntvE* group. The *Control* and *seaweed* groups had, on average, the lowest intake of the 2S stereoisomers, which was < 0.01 of their total α -tocopherol intake.

Compared to the combined mean of *Control* and *seaweed* groups, cows in the *NatvE* and *SyntvE* groups had 31 and 12% more ($P < 0.001$) plasma α -tocopherol, respectively (Table 4). Plasma concentration of the natural RRR isomer was 30% higher in the *NatvE* fed cows than the average of the other treatments, whilst cows fed *SyntvE* had on average eight times higher plasma

concentration of the synthetic isomers than cows on the other treatments. Almost all of the α -tocopherol in plasma was in the 2R stereoisomer form, and > 0.86 of the total was the RRR isomers on all treatments (Table 5). Despite 0.23 of total α -tocopherol intake being 2S, only *c.* 0.01 was in the 2S form in the plasma of the *SyntvE* group. Cows in the *Control*, *NatvE* and *seaweed* groups had plasma α -tocopherol stereoisomers from the synthetic derivative at very low yet measurable levels.

Milk and milk fat α -tocopherol concentration was affected ($P < 0.001$) by the treatments (Table 4). Cows in the *NatvE* group had 25% higher milk α -tocopherol concentration than the *Control*, while the α -tocopherol content in milk from cows fed *SyntvE* and seaweed did not differ significantly from that of the *Control*. As in plasma, > 0.85 of the α -tocopherol of milk was in the form of RRR on *SyntvE* diet and this value was observed to be > 0.97 on the other diets (Table 5).

Bioavailability of α -tocopherol and its stereoisomer ratios

The concentration of α -tocopherol in plasma and milk in cows offered *NatvE* was only slightly greater (1.09 and 1.06 times, respectively) than for cows fed *SyntvE* when adjusted to account for the differences in α -tocopherol intake (Table 6). However, the increase in plasma and milk α -tocopherol concentration relative (adjusted) to the control treatment was approximately twice as high on *NatvE* than on *SyntvE*. The concentration of α -tocopherol in plasma and milk per unit of 2R α -tocopherol consumed in cows fed *NatvE* supplement was 0.87 and 0.84 times that of cows fed *SyntvE* and was approximately the same (≈ 1) when calculated as the relative increase from the cows fed the control diet.

Antibody response to immunization

The summary of antibody titers (\log_2) is provided in Table 7. There was no effect of vitamin E supplementation on immunization response to the specific antigens with the exception of EQSA which showed a slight tendency ($P = 0.088$) to be lower under the *NatvE* group.

DISCUSSION

The present experiment was conducted to test the effects of feeding supplemental sources of vitamin E (RRR- α -tocopheryl acetate or all-rac- α -tocopheryl acetate) relative to seaweed and control diet on the vitamin E and its stereoisomers content of plasma and milk, and the adaptive immune response to immunization of mid-lactation dairy cows fed a basal feed of silage augmented with concentrates.

Plasma and milk α -tocopherol

The observed differences in α -tocopherol from concentrate intake were attributed to differences in the amount of concentrate DMI, whereas the difference from experimental supplements was due to the concentration of α -tocopherol in the supplements as designed. In addition to the *SyntvE* group, cows in the remaining three treatment groups yielded plasma and milk α -tocopherol stereoisomers of the *SyntvE* derivative at very low yet measurable levels. There are some possible reasons for this. Firstly, it could be due to the fact that a limited amount of the α -tocopherol stored in liver and other tissues from previous periods could be circulating at the background and hence available in the plasma and milk. A good example comes from the works of Weiser *et al* (1996), where control rats fed a vitamin E free diet for 4 months (pre-experimental period) and devoid of α -tocopherol for an extra 90 days (experimental period) showed traceable levels of α -tocopherol in

different tissues and plasma. Secondly, chemical analysis on the experimental supplements fed to the cows, except for the *seaweed* one, showed traceable levels of stereoisomers of the *SyntvE* derivatives suggesting that there might have been cross-contamination whilst sampling or feeding. However, the levels were very small and would not have altered the outcome of the treatments and, hence, our hypotheses. That said, the current study confirms that natural α -tocopherol is the predominating form in blood and milk of dairy cows, irrespective of dietary sources (Meglia *et al.* 2006; Weiss *et al.* 2009). The observed pattern of mean plasma and milk α -tocopherol concentration in the current study was in agreement with previous reports (Meglia *et al.* 2006; Weiss *et al.* 2009) where periparturient cows were fed either a control diet, supplemented with different forms of vitamin E and plasma concentration of α -tocopherol was measured. Indeed, in absolute terms, the values observed in the current paper are much higher than in previous works, but this was understandable considering the differences in terms of stage of lactation of the cows, type and quality of basal feed and also level of vitamin E supplementation.

Even though the intake of Vitamin E on the control diet was much lower than the recommended value, i.e. 2.6 IU/kg BW (NRC 2001) for late gestation and during lactation, the observed plasma α -tocopherol concentration is more than adequate. As such, the mean plasma α -tocopherol for control group (9.99 mg/l) was well above what is considered the minimum level (3–3.5 mg/l) to avoid health risks with periparturient dairy cows (Weiss *et al.* 1997). Furthermore, the value is also far higher than that reported for Swedish Holstein dairy cows in their mid-lactation and fed on organic rations (Lindqvist *et al.* 2011) and Norwegian dairy cows (6.9 mg/l) under winter feeding conditions (Sivertsen *et al.* 2005). Concentrations were, as per the hypothesis, lowest for cows in the control and seaweed group, intermediate for cows supplemented with *SyntvE*, and highest for cows fed supplemented with *NatvE*. This followed the trends of the concentrations in the diet consumed by the respective group.

Concentrations of α -tocopherol in plasma, milk and milk fat were about 1.2 times higher in the group supplemented with *NatvE* than in the *SyntvE* (Table 6). This agrees well with previous work with periparturient dairy cows (Meglia *et al.* 2006) using different sources of vitamin E. The milk α -tocopherol concentration falls a bit short of the results from Weiss *et al.* (2009) where the relative concentration was reported to be 1.24 to 1.43 times higher for cows supplemented with *NatvE* than for *SyntvE* with periparturient dairy cows. This could be due mainly to the elevated intake of α -tocopherol from basal feeds (silage and concentrates), in light of the observed values for the control group, where additional supplementation would have yielded less of an effect because of saturation.

Whilst comparing the *Control* group with the *seaweed* group, these two groups showed similar values from total α -tocopherol intake to its concentration in plasma and milk. However, there are other things which have not been looked into; for example, the incidence of mastitis, long term reproductive performance and milk iodine content. The latter, for example, was superior in seaweed meal fed group compared to standard mineral mixture fed Norwegian cows as reported by Jensen *et al.* (1968).

Plasma and milk α -tocopherol stereoisomers

The stereoisomer composition in milk followed similar pattern to those observed in plasma within each treatment. This is in agreement with other works on dairy cows and suggests either lack of preferential uptake of stereoisomers by mammary cells once in the blood (Meglia *et al.* 2006) or the limited availability of these stereoisomers in blood plasma (Jensen & Lauridsen 2007). Furthermore, in support of Meglia *et al.* (2006) and Weiss *et al.* (2009), the bioavailability of the RRR-stereoisomer in the current study was greater compared to the other stereoisomer forms of α -

tocopherol, irrespective of the source of supplementation, as indicated by the enrichment of the RRR stereoisomer in milk to > 0.86 of the total concentration.

Among the cows supplemented with *SyntvE* the relative enrichment of stereoisomers in plasma and milk was such that > 0.98 was in the 2R forms, whilst a very small proportion (*c.* 0.01–0.015) was in the 2S forms. Furthermore, within the 2R stereoisomers, the RRS, RSR and RSS stereoisomers were in a considerably lower concentration than the RRR form. The observed proportion of 2R vs. 2S was a big contrast to the proportion of these stereoisomers (0.77 and 0.23, respectively) in the feed fed to the cows. With periparturient cows fed daily with 917 mg all-rac- α -tocopheryl acetate on top of the intake from basal ration (300–525 mg RRR- α -tocopherol), Meglia *et al.* (2006) reported similar values. Likewise, Jensen *et al.* (2005) as described in Jensen & Lauridsen (2007), fed cows a daily dose of 3000 mg all-rac- α -tocopheryl acetate in addition to the 450 mg RRR- α -tocopherol occurring naturally in the basal ration for 16 days and reported comparable values in the plasma of cows. Furthermore, with sows fed on synthetic forms of vitamin E, Jensen & Lauridsen (2007) indicated that the 2R stereoisomers in total contributed *c.* 0.92–0.94 of the stereoisomers in milk whereas the 2S forms put together contributed the remaining small part (0.06–0.08). However, the same authors also reported differences in stereoisomer proportions in calf and lamb plasma compared to cows.

The preferential enrichment of the 2R form over the 2S form was suggested to be due to the action of hepatic α -tocopherol transport protein (α -TPP) that transfers liver α -tocopherol into plasma lipoproteins for extra-hepatic tissue delivery (Traber *et al.* 2008).

Responses to immunizations

Specific antibody titer values to immunization with EQSA antigen showed a tendency for dairy

cows in the *NatvE* group to be lower compared to other groups. This was unexpected but at the level of α -tocopherol intake and its plasma concentration observed in the current experiment, it can only be speculated that this could have been due to increased level of α -tocopherol radicals (Rietjens *et al.* 2002; Nwose *et al.* 2008). Such was the case with increased mastitis incidence in dairy cows that alluded to the absence of balance between vitamin E and scavenging molecules for vitamin E radicals (Bouwstra *et al.* 2010a, b). In human subjects with diabetes, Nwose *et al.* (2008) cautioned that vitamin E supplementation should be justified when the vitamin E regeneration system, which is responsible for balancing the outcome, is adequate. Rietjens *et al.* (2002) suggest that supplementation with vitamin E may not always exert the protective effect aimed for, as the effects may depend on the subtle redox equilibrium within the cells and the balance within the complete cellular antioxidant network.

The lack of anticipated difference of the effect of supplementation with seaweed on antibody titres warrants further investigation. However, a progression of specific antibody development has been observed in each group from pre-immunization values, as anticipated. The clear and elevated response to immunization suggests that the animals were at least not immunocompromised.

Conclusions

As hypothesized, supplementation of dairy cows with vitamin E, regardless of its form, increased total plasma and milk α -tocopherol compared to non-supplemented cows. This increment was bigger with *NatvE* than *SyntvE* supplementation. The lack of effects of vitamin E supplementation to elicit adaptive immune response against immunizations could be due to the fact that intake of vitamin E from the basal diets was not limiting. Therefore, based on our observation, it can be concluded that with good quality silage the need for supplemental vitamin E is not critical even

during the winter period with mid-lactation dairy cows.

We acknowledge the great help from the staff at the Agricultural Production Experimental Centre, Norwegian University of life Sciences. We thank Elisabeth Furuseth Hansen and Grethe Marie Johansen at the Norwegian School of Veterinary Science, Oslo, Norway for performing the laboratory analysis of the immunological parameters. We are grateful to Kristin Sørheim and Prof Knut Hove for taking the blood samples. We thank TINE SA for financial support of Ingvild Luteberget Nesheim's MSc work. Funding from The Foundation for Research Levy on Agricultural Products, the Agricultural Agreement Research Fund and the Research Council of Norway is much appreciated (Project number 190301/I10). The authors have no financial or other conflict of interest in the manuscript.

REFERENCES

- AL-MABRUK, R.M., BECK N.F.G. & DEWHURST, R.J. (2004). Effects of silage species and supplemental vitamin E on the oxidative stability of milk. *Journal of Dairy Science* **87**, 406-412.
- ALLEN, V. G., POND, K. R., SAKER, K. E., FONTENOT, J. P., BAGLEY, C. P., IVY, R. L., EVANS, R. R., BROWN, C. P., MILLER, M. F., MONTGOMERY, J. L., DETTLE, T. M. & WESTER, D. B. (2001). Tasco-Forage: III. Influence of a seaweed extract on performance, monocyte immune cell response, and carcass characteristics in feedlot-finished steers. *Journal of Animal Science* **79**, 1032-1040.
- BEECKMAN, A., VICCA, J., VAN RANST, G., JANSSENS, G. P. & FIEVEZ, V. (2010). Monitoring of vitamin E status of dry, early and mid-late lactating organic dairy cows fed conserved

- roughages during the indoor period and factors influencing forage vitamin E levels. *Journal of Animal Physiology and Animal Nutrition* (Berl) **94**, 736-746.
- BLIGH, E. G. & DYER, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology* **37**, 911-917.
- BOUWSTRA, R. J., GOSELINK, R. M. A., DOBBELAAR, P., NIELEN, M., NEWBOLD, J. R. & VAN WERVEN, T. (2008). The relationship between oxidative damage and vitamin e concentration in blood, milk, and liver tissue from vitamin E supplemented and nonsupplemented periparturient heifers. *Journal of Dairy Science* **91**, 977-987.
- BOUWSTRA, R. J., NIELEN, M., NEWBOLD, J. R., JANSEN, E. H. J. M, JELINEK, H. F. & VAN WERVEN, T. (2010a). Vitamin E supplementation during the dry period in dairy cattle. Part II: Oxidative stress following vitamin E supplementation may increase clinical mastitis incidence postpartum. *Journal of Dairy Science* **93**, 5696-5706.
- BOUWSTRA, R. J., NIELEN, M., STEGEMAN, J. A., DOBBELAAR, P., NEWBOLD, J. R., JANSEN, E. H. J. M & VAN WERVEN, T. (2010b). Vitamin E supplementation during the dry period in dairy cattle. Part I: Adverse effect on incidence of mastitis postpartum in a double-blind randomized field trial. *Journal of Dairy Science* **93**, 5684-5695.
- DERSJANT-LI, Y. & PEISKER, M. (2010). A critical review of methodologies used in determination of relative bio-availability ratio of *RRR*- α -tocopheryl acetate and *all-rac*- α -tocopheryl acetate. *Journal of the Science of Food & Agriculture* **90**, 1571-1577.
- EUROPEAN COMMISSION (1991). Council Regulation (EEC) No 2092/91 of 24 June 1991 on organic production of agricultural products and indications referring thereto on agricultural products and foodstuffs. *Official Journal of the European Communities* **L198**, 1-15
- EUROPEAN COMMISSION (1999). Council Regulation (EC) No 1804/1999 of 19 July 1999 supplementing Regulation (EEC) No 2092/91 on organic production of agricultural products

and indications referring thereto on agricultural products and foodstuffs to include livestock production. *Official Journal of the European Communities* **L222**, 1-28

EUROPEAN COMMISSION (2005). Council Regulation (EC) No 1567/2005 of 20 September 2005 amending Regulation (EEC) No 2092/91 on organic production of agricultural products and indications referring thereto on agricultural products and foodstuffs. *Official Journal of the European Communities* **L252**, 1-1

EUROPEAN COMMISSION (2009). Commission regulation (EC) No 152/2009 of 27 January 2009 laying down the methods of sampling and analysis for the official control of feed. *Official Journal of the European Communities* **L54**, 1-130.

GOVASMAR, E., STEEN, A., STRØM, T., HANSEN, S., SINGH, B. R. & BERNHOFT, A. (2005). Status of selenium and vitamin E on Norwegian organic sheep and dairy cattle farms. *Acta Agriculturae Scandinavica, Section A – Animal Science* **55**, 40-46.

IFOAM (2007). *The IFOAM Norms for Organic Production and Processing, version 2005*. Bonn, Germany: IFOAM.

JENSEN, S.K. (2008). Improved Bligh and Dyer extraction procedure. *Lipid Technology* **20**, 280–281.

JENSEN, A., NEBB, H. & SÆTER, E. A. (1968). *The Value of Norwegian seaweed Meal, as a Mineral Supplement for Dairy Cows*. Oslo: Norwegian Institute of seaweed Research.

JENSEN, S. K., JENSEN, C., JAKOBSEN, K., ENGBERG, R. M., ANDERSEN, J. O., LAURIDSEN, C., SØRENSEN, P., SKIBSTED, L. H. & BERTELSEN, G. (1998). Supplementation of broiler diets with retinol acetate, β -carotene or canthaxanthin: Effect on vitamin status and oxidative status of broilers *in vivo* and on meat stability. *Acta Agriculturae Scandinavica, Section A – Animal Science* **48**, 28-37.

JENSEN, S. K., KRISTENSEN, N. B., LAURIDSEN, C. & SEJERSEN, K. (2005). Enrichment of cows'

milk with natural or synthetic vitamin E. In *Proceedings of the 10th Symposium on Vitamins and Additives in Nutrition of Man and Animal 28-29 September, Jena, Thuringia, Germany* (Eds R. Schubert, G. Flachowsky, G. Jahreis & R. Bitsch), pp. 78-83. Braunschweig, Germany: Bundesforschungsanstalt für Landwirtschaft (FAL).

JENSEN, S. K. & LAURIDSEN, C. (2007). α -Tocopherol stereoisomers. *Vitamins & Hormones* **76**, 281-308.

JENSEN, S. K. & NIELSEN, K. N. (1996). Tocopherols, retinol, beta-carotene and fatty acids in fat globule membrane and fat globule core in cows' milk. *Journal of Dairy Research* **63**, 565-574.

LEE, S.K., KIM, Y.S., LIANG, C.Y. & SONG, Y.H. (2003). Effects of dietary vitamin E supplementation on color stability, lipid oxidation and reducing ability of Hanwoo (Korean cattle) beef during retail display. *Asian-Australasian Journal of Animal Sciences* **16**, 1529-1534.

LINDQVIST, H., NADEAU, E., PERSSON WALLER, K., JENSEN, S. K. & JOHANSSON, B. (2011). Effects of RRR- α -tocopheryl acetate supplementation during the transition period on vitamin status in blood and milk of organic dairy cows during lactation. *Livestock Science* **142**, 155-163.

LIU, Q., LANARI, M. C. & SCHAEFER, D. M. (1995). A review of dietary vitamin E supplementation for improvement of beef quality. *Journal of Animal Science* **73**, 3131-3140.

MEGLIA, G. E., JENSEN, S. K., LAURIDSEN, C. & WALLER, K. P. (2006). Alpha-tocopherol concentration and stereoisomer composition in plasma and milk from dairy cows fed natural or synthetic vitamin E around calving. *Journal of Dairy Research* **73**, 227-234.

MERTENS, D. R. (2002). Gravimetric determination of amylase-treated neutral detergent fiber in feeds with reflusing in beakers or crucibles: collaborative study. *Journal of AOAC International* **85**, 1217-1240.

MOGENSEN, L., KRISTENSEN, T., SØEGAARD, K., JENSEN, S. K. & SEHESTED, J. (2012). Alfa-

tocopherol and beta-carotene in roughages and milk in organic dairy herds. *Livestock Science* **145**, 44-54.

NRC (2001). *Nutrient Requirements of Dairy Cattle*. Seventh Revised Edition ed. Washington, D.C.: National Academy Press.

NWOSE, E. U., JELINEK, H. F., RICHARDS, R. S. & KERR, P. G. (2008). The 'vitamin E regeneration system' (VERS) and an algorithm to justify antioxidant supplementation in diabetes - A hypothesis. *Medical Hypotheses* **70**, 1002-1008.

OTTERLEI, M., OSTGAARD, K., SKJAAK-BRAEK, G., SMIDSRØD, O., SOON-SHIONG, P. & ESPEVIK, T. (1991). Induction of cytokine production from human monocytes stimulated with alginate. *Journal of Immunotherapy* **10**, 286-291.

RIETJENS, I. M. C. M., BOERSMA, M. G., DE HAAN, L., SPENKELINK, B., AWAD, H. M., CNUBBEN, N. H., VAN ZANDEN, J. J., VAN DER WOUDE, H., ALINK, G. M. & KOEMAN, J. H. (2002). The pro-oxidant chemistry of the natural antioxidants vitamin C, vitamin E, carotenoids and flavonoids. *Environmental Toxicology and Pharmacology* **11**, 321-333.

ROKSTAD, A. M., ESPEVIK, T., SKJÅK-BRÆK, G. & VADSTEIN, O. (1996). Poly mannuronic acid (poly M) and lipopolysaccharide (LPS) induced superoxide anion production in Atlantic salmon (*Salmo salar L.*) macrophages and the dependence of serum. In *Fish Vaccinology: Developments in Biological Standardization* (Eds F. Brown, R. Gudding, A. Lillehaug & Paul J. Midtlyng), p. 434. Basel, Switzerland: S. Karger.

SIVERTSEN, T., OVERNES, G. OSTERAS, O., NYMOEN, U. & LUNDER, T. (2005). Plasma vitamin E and blood selenium concentrations in Norwegian dairy cows: Regional differences and relations to feeding and health. *Acta Veterinaria Scandinavica* **46**, 177-191.

SKJERMO, J. & BERGH, Ø. (2004). High-M alginate immunostimulation of Atlantic halibut (*Hippoglossus hippoglossus L.*) larvae using *Artemia* for delivery, increases resistance against

vibriosis. *Aquaculture* **238**, 107-113.

TRABER, M. G., FREI, B. & BECKMAN, J. S. (2008). Vitamin E revisited: do new data validate benefits for chronic disease prevention? *Current Opinion in Lipidology* **19**, 30-38.

WEISER, H., RISS, G. & KORMANN, A. W. (1996). Biodiscrimination of the eight *alpha*-tocopherol stereoisomers results in preferential accumulation of the four 2r forms in tissues and plasma of rats. *Journal of Nutrition* **126**, 2539-2549.

WEISS, W. P., HOGAN, J. S., TODHUNTER, D. A. & SMITH, K. L. (1997). Effect of vitamin E supplementation in diets with a low concentration of selenium on mammary gland health of dairy cows. *Journal of Dairy Science* **80**, 1728-1737.

WEISS, W. P., HOGAN, J. S. & WYATT, D. J. (2009). Relative bioavailability of all-rac and RRR vitamin E based on neutrophil function and total α -tocopherol and isomer concentrations in periparturient dairy cows and their calves. *Journal of Dairy Science* **92**, 720-731.

Table 1. *Ingredient composition of the four experimental supplements prepared with no added vitamin E (Control), RRR- α -tocopheryl acetate (NatvE), all-rac- α -tocopheryl acetate (SyntvE) and seaweed and fed to cows in their mid-lactation as a source of α -tocopherol on top of the basal feed*

Ingredient (in mg/kg, unless stated)	Control	NatvE	SyntvE	seaweed
Barley (g/kg)	749	740	744	515
seaweed meal (g/kg)	0	0	0	305
Molasses (g/kg)	42	42	42	36
Vitamin E*	0	12589	7591	0
Calcium carbonate (CaCO ₃) (g/kg)	52.1	51.4	51.6	40.7
Monocalcium phosphate (Ca(H ₂ PO ₄) ₂ .H ₂ O) (g/kg)	15.6	15.3	15.4	15.9
Magnesium phosphate (MgHPO ₄ .H ₂ O) (g/kg)	75.9	74.8	75.2	59.4
Sodium chloride (NaCl) (g/kg)	53.3	52.6	52.8	23.3
Sodium selenite (Na ₂ SeO ₃)	153	150	151	130
Zinc sulphate (ZnSO ₄ .H ₂ O)	2633	2596	2608	2200
Manganese sulphate (MnSO ₄ .H ₂ O)	2153	2122	2132	1810
Calcium iodate (Ca(IO ₃) ₂)	54	54	54	0
Cobalt carbonate (2CoCO ₃ .3Co(OH) ₂ .H ₂ O)	46	45	46	39
Copper sulphate (CuSO ₄)	836	824	828	0
Vitamin A (retinyl acetate)	184	181	182	0
Vitamin D ₃ (cholecalciferol)	54	54	54	0

*SyntvE: 500000 mg All-rac- α -tocopherylacetate/kg vitamin E-product, i.e. 3800 mg α -tocopherylacetate/kg supplement

RRR: 301000 mg RRR- α -tocopherylacetate/kg vitamin E-product, i.e. 3800 mg α -tocopherylacetate/kg supplement

Table 2. Chemical composition of the four experimental supplements [no added vitamin E (Control), RRR- α -tocopheryl acetate (NatvE), all-rac- α -tocopheryl acetate (SyntvE) and seaweed] and basal feeds (silage and concentrate), and proportions (of total) of α -tocopherol stereoisomers. Values in brackets are standard deviations (n=4)

Item	Experimental supplements				Basal feeds	
	Control	NatvE	SyntvE	seaweed	Silage	Concentrate
DM (g/kg)	880 (3.5)	892 (4.6)	879 (5.4)	865 (5.4)	283 (1.5)	875 (2.4)
CP (g/kg DM)	86 (1.0)	85 (2.0)	88 (1.2)	89 (1.2)	123 (3.9)	163 (4.1)
aNDFom (g/kg DM)	137 (3.8)	133 (7.9)	137 (1.8)	228 (5.4)	487(12.0)	287 (5.7)
Fat (g/kg DM)	29 (3.1)	28 (1.6)	31 (6.3)	27 (2.1)	52 (3.7)	81 (3.4)
<u>Silage fermentation products (g/kg DM)</u>						
Ammonia N (g/kg N)					53 (5.5)	
Ethanol					6 (2.3)	
Lactic acid					46 (3.6)	
Volatile fatty acids					15 (1.3)	
<u>α-Tocopherol</u>						
Total (mg/kg DM)	79.3(10.3)	1052 (243)	930 (227)	4.4 (1.8)	39.0 (2.4)	7.9 (0.50)
RRR	0.73 (0.054)	0.89 (0.013)	0.12 (0.003)	ND	1.00 (0.0)	ND
RSS	0.03 (0.011)	0.005 (0.001)	0.13 (0.003)	ND	0 (0.0)	ND
RRS	0.04 (0.020)	0.04 (0.005)	0.13 (0.002)	ND	0 (0.0)	ND
RSR	0.03 (0.010)	0.006 (0.002)	0.12 (0.005)	ND	0 (0.0)	ND
2S	0.18 (0.026)	0.06 (0.007)	0.51 (0.009)	ND	0 (0.0)	ND
<u>Fatty acids (g/kg DM)</u>						
C14:0	0.1 (0.0)	0.1 (0.0)	0.1 (0.01)	1.1 (0.04)	0.2 (0.04)	0.1 (0.0)
C16:0	4.9 (0.11)	4.9 (0.07)	5.0 (0.08)	4.3 (0.05)	2.7 (0.05)	9.9 (0.07)
C18:0	0.2 (0.00)	0.2 (0.00)	0.3 (0.01)	0.3 (0.01)	0.3 (0.02)	1.45 (0.04)
c9-C18:1	2.0 (0.04)	1.9 (0.05)	2.1 (0.06)	5.3 (0.16)	0.7 (0.07)	28.4 (1.50)
C18:2n6	6.5 (0.38)	5.8 (0.38)	6.3 (0.23)	6.6 (0.12)	3.3 (0.10)	25.1 (0.36)
C18:3n3	0.5 (0.04)	0.5 (0.03)	0.5 (0.03)	0.9 (0.01)	8.8 (0.14)	2.1 (0.19)
Total	14.7(0.60)	14.0 (0.48)	14.9 (0.29)	21.1 (0.35)	16.9 (0.11)	70.0 (2.27)

ND: Not detectable

Table 3. Daily DM and α -tocopherol intake by dairy cows in their mid-lactation fed one of the four different experimental supplements: no added vitamin E (Control), RRR- α -tocopheryl acetate (NatvE), all-rac- α -tocopheryl acetate (SyntvE) and seaweed (n=24)

Description	Treatment group				S.E.M.	D.F.	P
	Control	NatvE	SyntvE	seaweed			
<u>DM intake (kg/d)</u>							
Silage	14.6	14.7	14.6	14.4	0.51	7.9	NS
Concentrate	2.92	2.87	2.86	2.99	0.047	11.8	0.021
Experimental supplements	0.55	0.52	0.51	0.63	0.016	-	-
Total intake	18.1	18.1	18.0	18.1	0.47	8.4	NS
Refuse	0.17	0.24	0.26	0.06	0.063	31.9	0.001
<u>α-tocopherol intake (mg/d) by source</u>							
Silage	570	574	571	566	19.9	7.9	NS
Concentrate	23.0	22.7	22.6	23.6	0.38	32.8	0.004
Experimental supplements	43.5	549	477	2.8	13.48	19.4	<0.001
Total	636	1146	1070	590	16.8	49.4	<0.001
<u>α-tocopherol intake (mg/d) by stereoisomer category</u>							
RRR	624	1086	649	591	16.3	13.3	<0.001
Sum other 2R (RSS, RRS, RSR)	4	28	117	0	2.8	27.5	<0.001
Sum 2S	8	32	244	0	3.8	26.9	<0.001
<u>Vitamin E intake</u>							
Total IU/d	948	1707	1360	880	24.1	50.5	<0.001
IU/kg LW	1.71	3.02	2.43	1.56	0.055	39.8	<0.001

D.F. = Satterthwaite's approximation for degrees of freedom

S.E.M. = Standard error of the mean

NS, not significant

IU of vitamin E was calculated as 1.49 x mg of α -tocopherol from feedstuffs and for NatvE supplement and 1 x mg of α -tocopherol for the SyntvE supplement

LW = Live weight

Table 4. Daily milk yield, milk fat content, milk and plasma α -tocopherol concentration in dairy cows in their mid-lactation ($n = 24$) fed daily supplements of no added vitamin E (Control), RRR- α -tocopheryl acetate (NatvE), all-rac- α -tocopheryl acetate (SyntvE) and seaweed

Description	Treatment group				S.E.M.	D.F.	P
	Control	NatvE	SyntvE	seaweed			
Milk yield (kg/d)	15.7	15.9	16.3	16.0	0.64	2.4	NS
Milk fat (g/kg)	42.7	42.5	41.1	41.2	1.23	6.6	0.047
<i><u>α-Tocopherol in plasma (mg/l)</u></i>							
Total	9.99	13.17	11.27	10.13	0.593	8.2	<0.001
RRR	9.83	12.82	9.69	10.05	0.544	8.5	<0.001
Sum other 2R	0.15	0.33	1.47	0.07	0.076	22.0	<0.001
Sum 2S	0.01	0.02	0.11	0.01	0.005	78.7	<0.001
<i><u>α-Tocopherol in milk</u></i>							
Total (mg/kg fat)	28.0	35.3	30.5	27.6	1.18	29.3	<0.001
Total (mg/l)	1.20	1.50	1.25	1.14	0.052	35.4	<0.001
RRR (mg/l)	1.20	1.47	1.09	1.14	0.050	33.9	<0.001
Sum other 2R (mg/l)	0.00	0.03	0.14	0.00	0.008	23.2	<0.001
Sum 2S (mg/l)	0.001	0.005	0.019	0.001	0.0011	77.9	<0.001

D.F. = Satterthwaite's approximation for degrees of freedom

S.E.M. = Standard error of the mean

NS, not significant

Table 5. Relative proportions of α -tocopherol stereoisomers in the diets, plasma and milk samples from dairy cows in their mid-lactation fed on no added vitamin E (Control), RRR- α -tocopheryl acetate (NatvE), all-rac- α -tocopheryl acetate (SyntvE) and seaweed

Description	Treatment group				S.E.M	D.F.	P- value
	Control	NatvE	SyntvE	seaweed			
<u>Diet</u>							
RRR	0.981	0.947	0.607	1.000	0.0063	14.8	<0.001
RSS	0.002	0.002	0.056	0.000	0.0008	17.0	<0.001
RRS	0.002	0.019	0.056	0.000	0.0010	12.3	<0.001
RSR	0.002	0.003	0.054	0.000	0.0008	16.8	<0.001
Sum 2S	0.012	0.028	0.228	0.000	0.0036	14.8	<0.001
<u>Plasma</u>							
RRR	0.984	0.974	0.866	0.992	0.0045	76.3	<0.001
RSS	0.004	0.008	0.041	0.002	0.0014	40.3	<0.001
RRS	0.007	0.011	0.048	0.003	0.0017	76.0	<0.001
RSR	0.004	0.006	0.036	0.002	0.0013	27.9	<0.001
Sum 2S	0.001	0.001	0.010	0.001	0.0004	76.8	<0.001
<u>Milk</u>							
RRR	0.994	0.979	0.874	0.996	0.0060	23.0	<0.001
RSS	0.001	0.004	0.034	0.001	0.0015	26.9	<0.001
RRS	0.002	0.007	0.039	0.001	0.0021	26.5	<0.001
RSR	0.002	0.007	0.038	0.001	0.0021	20.3	<0.001
Sum 2S	0.001	0.003	0.015	0.001	0.0008	71.0	<0.001

D.F. = Satterthwaite's approximation for degrees of freedom

S.E.M. = Standard error of the mean

NS, not significant

Table 6. Ratios of relative intakes of α -tocopherol and relative concentrations of α -tocopherol in plasma and milk from cows fed *NatvE* or *SyntvE* supplement

Ratios	<i>NatvE/SyntvE</i> *	Intake adjusted <i>NatvE/SyntvE</i>	
		α -tocopherol [†]	2R [‡]
Intakes			
α -Tocopherol, total	1.07	-	-
2R Tocopherol	1.35	-	-
RRR Tocopherol	1.67		
<i>Intake incremental relative to:</i>			
Control, total	1.18	-	-
seaweed, total	1.16	-	-
Control, 2R	2.45	-	-
Plasma α -tocopherol	1.17	1.09	0.87
Plasma α -tocopherol increment relative to value from control group	2.48	2.11	1.01
Milk α -tocopherol	1.14	1.06	0.84
Milk α -tocopherol increment relative to value from control group	2.33	1.98	0.95

**NatvE/SyntvE* ratio = mean intake, concentration and increments relative to control and seaweed for *NatvE* treatment divided by mean of *SyntvE* treatment (see Tables 3 and 4 for original data).

[†] [(α -tocopherol concentration in *NatvE*/ α -tocopherol concentration in *SyntvE* treatment)/(α -tocopherol intake by *NatvE*/ α -tocopherol intake by *SyntvE* treatment)].

[‡] [(α -tocopherol concentration in *NatvE*/ α -tocopherol concentration in *SyntvE* treatment)/(2R α -tocopherol intake by *NatvE* /2R α -tocopherol intake by *SyntvE* treatment)].

Table 7. Antibody titers (log₂) to human serum albumin (HSA), ovalbumin (OVA), equine serum albumin (EQSA) and canine serum albumin (CSA) from dairy cows (n=6) fed on basal feeds of silage and concentrate, and supplemented with no added vitamin E (Control), RRR- α -tocopheryl acetate (NatvE), all-rac- α -tocopheryl acetate (SyntvE) and seaweed

	<i>Treatment group</i>				S.E.M.	D.F.	<i>P</i>
	<i>Control</i>	<i>NatvE</i>	<i>SyntvE</i>	<i>seaweed</i>			
<u>Pre-vaccination values (SD)</u>							
HSA	9.6 (0.52)	9.5 (0.41)	9.8 (0.84)	9.5 (0.41)			
OVA	9.6 (0.82)	9.3 (0.0)	9.3 (0.0)	10.0 (1.21)			
EQSA	9.5 (0.0)	9.3 (0.0)	9.3 (0.0)	9.3 (0.0)			
CSA	9.3 (0.0)	9.3 (0.0)	9.3 (0.0)	9.3 (0.0)			
<u>Response to immunization</u>							
HSA	12.8	12.5	13.1	12.6	0.35	18.8	NS
OVA	12.6	12.1	12.3	12.8	0.25	20.0	NS
EQSA	13.5	12.3	13.1	13.0	0.30	20.0	0.086
CSA	12.1	11.0	11.8	11.3	0.36	19.5	NS

D.F. = Satterthwaite's approximation for degrees of freedom

S.E.M. = Standard error of the mean

NS, not significant