

Investigating large-scale brain dynamics

Investigating large-scale brain dynamics using field potential recordings: Analysis and interpretation

Bijan Pesaran^{1,2}, Martin Vinck³, Gaute T. Einevoll^{4,5}, Anton Sirota⁶, Pascal Fries^{3,7}, Markus Siegel⁸, Wilson Truccolo^{9,10}, Charles E. Schroeder^{11,12}, Ramesh Srinivasan¹³

¹Center for Neural Science
New York University
New York, NY USA

²NYU Neuroscience Institute
New York University Langone Health
New York, NY USA

³Ernst Strüngmann Institute (ESI) for Neuroscience in Cooperation with Max Planck Society
Frankfurt, Germany

⁴Faculty of Science and Technology
Norwegian University of Life Sciences
Ås, Norway

⁵Department of Physics
University of Oslo
Oslo, Norway

⁶Bernstein Center for Computational Neuroscience Munich
Munich Cluster of Systems Neurology (SyNergy)
Faculty of Medicine
Ludwig-Maximilians-Universität München
Planegg-Martinsried, Germany

⁷Donders Institute for Brain, Cognition, and Behaviour
Radboud University Nijmegen
Nijmegen, Netherlands

⁸Centre for Integrative Neuroscience & MEG Center
University of Tübingen
Tübingen, Germany

⁹Department of Neuroscience and Institute for Brain Science
Brown University
Providence, RI USA

¹⁰Center for Neurorestoration and Neurotechnology
U.S. Department of Veterans Affairs
Providence, RI, USA

¹¹Translational Neuroscience Division
Center for Biomedical Imaging and Neuromodulation
Nathan Kline Institute
Orangeburg, NY, USA

¹²Department of Neurosurgery

Investigating large-scale brain dynamics

Columbia College of Physicians and Surgeons
New York, NY, USA

¹³Department of Cognitive Sciences
Department of Biomedical Engineering
University of California
Irvine, CA USA

Word Count: 8000 words max

Abstract		164
A – Introduction	~500 words	549
B –Modeling	~1000 words	1126
C – Analyses and Interpretations	~5000 words	142
C1 – Activation		1243
C2 – Correlation		1306
C3 – Communication		793
C4 – Coding		821
D – Conclusion	~500 words	382
Box 1 - Biophysics of extracellular potentials		689
Box 2 - Interpreting local and remote sources		803
Figures		8
References		169 references
Extended Supplement		
S1 – Activation		
S1A - Spectral estimation		1691 words
S1B - Higher-order structure		446
S1C - Inverse modeling		816
S2 – Correlation		
S2A – Mathematical model of coherence		402
S2B – Spike-field coherence		289
S2C - Volume conduction		432
S2D - Experimental considerations		1747
S3 - Communication		
S3A - Time-delays		683
Supplementary References		109 references

Abstract

New technologies to record electrical activity from the brain on a massive scale offer tremendous opportunities for discovery. Electrical measurements of large-scale brain dynamics, termed field potentials, are especially important to understanding and treating the human brain. Here, our goal is to provide best practices on how field potential recordings (EEG, MEG, ECoG and LFP) can be analyzed to identify large-scale brain dynamics, and to highlight critical issues and limitations of interpretation in current work. We focus our discussion of analyses around the broad themes of activation, correlation, communication and coding. We provide recommendations for interpreting the data using forward and inverse models. The forward model describes how field potentials are generated by the activity of populations of neurons. The inverse model describes how to infer the activity of populations of neurons from field potential recordings. A recurring theme is the challenge of understanding how field potentials reflect neuronal population activity given the complexity of the underlying brain systems.

Part A – Introduction

Brain dynamics are generated by the activity of diverse and massive populations of interconnected neurons distributed across the nervous system. Complete system-wide recordings of neuronal population activity at the cellular scale would provide the ideal basis for analyzing brain dynamics. Unfortunately, tools to directly study the mapping from activity to dynamics at the cellular scale remain out of reach, especially in humans. Fortunately, neuronal dynamics are redundant and correlated. As a result, we can investigate brain dynamics by analyzing the spatiotemporal signals generated by populations of neurons that take the form of field potentials¹. Field potentials encompass a range of signals including local field potentials (LFP), electrocorticography (ECoG), intracranial stereotactic electroencephalography (stereo-EEG), scalp electroencephalography (EEG) and magnetoencephalography (MEG; **Fig 1**). All field potential recordings have good temporal precision in the millisecond range and are available in both humans and non-human animals. The techniques differ primarily in their spatial resolution, their coverage, and their degree of invasiveness.

Understanding how field potentials measure particular patterns of neuronal activity depends on a forward model and an inverse model. Forward models describe how the recorded potentials are generated by neuronal activity. Inverse models are used to analyze the recorded potentials and infer the underlying neuronal sources. The ingredients to both the forward and inverse model are conservation of charge and Maxwell's equations^{2,3}, electrical properties of brain and head tissues as well as the physics and geometry of the neural source, anatomy, and recording sensor. Forward models can, in principle, be used to precisely compute the recorded potentials. Inverse models, however, cannot in general compute the cellular patterns of activity from recorded potentials. This is because the problem is ill-posed and different patterns of neuronal activity can generate the same field potential measurements. Even with an accurate inverse model, we cannot reconstruct the exact pattern of cellular sources that generates the measurement.

Several aspects of field potential research have been previously reviewed⁴⁻⁹. The goal of the current paper is to guide advanced, but not expert, researchers in the analysis and interpretation of large-scale brain dynamics using field potential recordings. As we will demonstrate, there are many benefits to analyzing field potential recordings to study large-scale brain dynamics. The principal risks stem from the lack of a well-posed inverse problem. In **Part B - Modeling**, we outline field potential research and highlight some areas of recent controversy. This is supplemented by **Box 1 - Biophysics of extracellular potentials**. **Part C – Analyses and Interpretations** forms our main focus and centers on the appropriate use of the forward model. For this section, we adopt a didactic approach and present clear guidelines for best practices in analysis and interpretation of field potential recordings. We organize the material in four sections - **Activation, Correlation, Communication** and **Coding**. For some issues, we can point to solutions, such as how to detect correlations between different neuronal signals, and how to decode information present in neuronal activity. Other topics, such as how to disentangle local activity from activity in synaptic inputs from remote sources and how to infer causal influences are more complex and merit detailed study in **Box 2 – Interpreting local and remote sources**. An **Extended Supplement** provides the interested reader with more detailed supporting information and discussion.

Part B – Modeling

All extracellular potentials stem from summed contributions of transmembrane currents across the surfaces of neurons (and in principle also glia cells, see **Box 1**)^{4,5,10}. In this sense, the biophysical forward model of field potential recordings is similar to that of spike recordings. However, field potentials differ from single neuron recordings in important ways. Single neuron spike recordings have a single source –a neuron that generates action potentials at discrete events in time. The single unit measurement is a classification – we want to determine whether the neuron has generated an action potential. Distant sources can only contribute to the noise floor above which single neuron activity must be discriminated. In contrast, field potentials are continuous signals that do not have a single source⁵. A field potential measurement has contributions from different sources weighted according to the measurement modality – LFP, ECoG, EEG or MEG. Of the various potential sources, the largest cellular contribution comes from neurons that generate the largest current dipoles, those

Investigating large-scale brain dynamics

with extended, oriented dendritic arbors (see **Fig 2a**). Temporally correlated synaptic inputs at restricted dendritic sites also contribute strongly to field potentials. Distant sources can contribute directly to field potentials due to volume conduction or, for MEG, field spread. Volume conduction occurs because electromagnetic fields propagate through biological tissues^{6,11}. Consequently, field potentials reflect the organization of large-scale brain dynamics.

While biophysical forward-modeling schemes for field potential signals are well-established^{6,10,12}, several issues remain controversial. One controversial issue concerns the frequency-dependence of the electrical conductivity of the extracellular space, i.e., whether volume conduction biases LFP and EEG recordings towards certain frequencies¹³. Field potential power has been observed to decay in power-law like way ($1/f^\alpha$) for higher frequencies. Establishing a power law is challenging because the range of frequencies is limited, typically only spanning two or three orders of magnitude. While the reason for any power law remains unclear, potential explanations include scale-free dynamics, the capacitive effects of neuronal membranes, and noise from evenly distributed ion channels across dendrites^{14–16}. Early experiments found very little frequency dependence in the frequency range of interest for neuronal recordings^{17,18}, implying that the extracellular medium is essentially resistive, i.e., Ohmic with a negligible capacitive component. Subsequent observations showed a strongly reduced electrical conductivity for frequencies below 100 Hz¹⁹ suggesting a strong low-pass filtering of the LFP and EEG from the extracellular medium²⁰. A series of further studies, using a variety of measurement setups, have however confirmed the early findings and found only weak frequency dependencies^{21–25}. Thus, on balance the experimental observations seem to point to a largely resistive extracellular medium.

The contribution of spiking events to field potentials is somewhat controversial. In cortex and hippocampus, the high-frequency part of the extracellular potential, i.e., above some hundred Hz, is thought to be dominated by spiking activity. Likewise, the low-frequency part, i.e., below some tens of Hz is thought to be dominated by the sub-threshold extracellular signatures of synaptic activation^{4,5} although spikes also contribute²⁶. Spikes are far more likely to contribute to intracortical microelectrode LFPs than to the EEG recorded from a scalp electrode. The crossover frequency, i.e., the frequency above which the spike contributions start to dominate, depends on the competition between sub-threshold and spiking signal contributions. The result depends on brain area and brain state: In hippocampus, spikes contribute to the extracellular signal for frequencies down to 100 Hz^{27,28}. In monkey visual cortex, extracellular signal frequencies as low as ~50 Hz have been associated with spiking²⁹. In general, the contribution of spiking events depends on the shape and amplitude of the action potential waveform and the firing rate³⁰.

Inverse modeling to infer neuronal sources also presents controversial issues. Inferring neuronal sources from M/EEG is called source reconstruction. Direct validation has been performed with electrical stimulation of an implanted electrode^{31,32} and expected results have been obtained in the case of, for example, activity in early sensory cortices^{33–35} and in the case of dipole sources for epilepsy³⁶. However, in general, additional information is always required in addition to the recorded potentials or fields, sources may be distributed with many dipoles and/or higher-order multipoles and reconstruction depends on information that is often unavailable^{37,38}. Inferring neuronal sources from LFPs has typically involved current source density (CSD) analysis. Direct empirical validation of CSD has also proven challenging³⁷. Although the assumption of charge conservation is well established³, it can be difficult in practice to determine exactly where the currents flow.

Finally, the ability to read-out, or decode, information processing from the brain underlies the development of brain-machine interfaces and neural prostheses and depends on the choice of neuronal signal (spiking/LFP/ECOG/EEG/MEG). Field potential decoding analyses depend on how field potentials measure selectivity of the underlying neuronal sources as well as the spatiotemporal organization of the source selectivity. We can summarize these factors in a functional variant of the biophysical forward model. The functional forward model describes how neuronal activity varies due to different task conditions and how this generates selective field potential responses (see **Fig 2b**). This model involves the biophysical model, along with how populations of neurons encode behavioral or task-relevant variables. The functional forward model predicts that field potentials are more selective for field potential modalities in which the measurement projection is spatially-confined, at sites where the weighted sources encode similar variables, and for task variables that change the spatial-temporal activity correlations to which field potentials are sensitive.

Relatively little work has investigated field potential selectivity in terms of the forward model and the nature of field potential selectivity remains controversial. An early view was that LFP selectivity differed from spiking recordings and was similar to EEG selectivity. However, empirical studies show LFPs contain more sensory and movement information than EEG, and are more comparable to the information available in the spiking activity of single neurons^{39–45}. Information about reach, grasp and speech actions are also present in human ECoG recordings^{46,47}. While more work is needed, in cortex, the presence of field potential selectivity comparable to single neuron selectivity may reflect the underlying columnar organization of cortex^{39,41,44,45}. Since field potential recordings are easier to obtain than recordings of spiking activity, these and other studies suggest that ECoG and/or LFP signals, in addition to spiking, may serve as a basis for a high-performance brain-machine interface⁴⁸. For M/EEG, multivariate classification approaches also reveal neuronal encoding of various forms of sensory and cognitive information^{49–52}. M/EEG selectivity is also shaped by volume conduction in a process called spatial filtering^{6,53}. If many sources with different dynamics occur in dipole layers of different sizes, the measured M/EEG signal is dominated by the temporal dynamics of those source components that are synchronized more broadly over the cortical surface^{5,37}. Consistent with these observations, M/EEG signals appear to be less selective than ECoG signals which appear to be less selective than LFP signals^{39,54,55}.

C – Analyses and Interpretations

Field potential recordings are time series – a series of measurements ordered in time – that reflect neuronal dynamics. As a result, time series analysis tools that measure dynamics, such as spectral analysis⁵⁶, are often used to characterize field potentials^{57,58}. The first three sections present time series analysis and interpretations that depend on the biophysical forward model. In **C1 – Activation**, we discuss how to assess the signal characteristics at a given site and their changes with experimental conditions. In **C2 - Correlations** we discuss how to characterize correlations in activity between sites. In **C3 – Communication**, we detail tests for directed influences that may reflect neuronal communication. **C4 – Coding**, considers the functional aspects of the forward model. We discuss neuronal coding and how information in the underlying neuronal activity can be decoded from field potentials. In each section, we discuss the associated analyses and interpretations.

C1 - Activation

We first focus on estimating the magnitude of the activity and testing for differences, before turning to interpreting spectral features and oscillations.

Statistical estimation: Two classic measures of time series are the autocorrelation or autocovariance and the power spectral density function, which we will refer to as the spectrum (**Fig 3**). The autocorrelation measures how much activity is correlated at two points separated in time, $\tau = t_2 - t_1$, also called lag. The spectrum measures how much power the activity contains at each frequency, f . The spectrum and autocorrelation function provide complementary information but are not equivalent measures of activity. The spectrum reveals spectral peaks (that may correspond to physical sources) that can be difficult to distinguish in the autocorrelation function. Conversely, the autocorrelation function reveals how rapidly a signal tends to become uncorrelated over time, which is not directly evident from the spectrum itself.

Spectrum and correlation function estimates assume that neuronal activity is stationary. Stationarity means the signal statistics, power, correlations, remain constant over time. Strictly-speaking, stationarity is never satisfied. Non-stationarities occur with transient events. The most extreme example is the non-stationarity of activity around the time of stimulus onset. However, even in the absence of changes in sensory input, brain activity displays transient events that can wax and wane, such as beta activity in the frontal cortices and ripples in the hippocampus. Non-stationarities can also occur across different time scales, such as fluctuations in high-frequency synchronization following low-frequency excitability changes⁵⁹, and can occur when activity appears stationary, e.g. sustained stimulus-induced gamma oscillations in visual cortex, sustained theta oscillations in rodent hippocampus during walking. Changes also occur spontaneously, for example as one spontaneously

Investigating large-scale brain dynamics

drifts back and forth from an aroused to a non-aroused state^{60,61}. In each case, the spectrum/auto-correlation averaged across these fluctuations will not be a good descriptor of the data.

If the changes occur relatively slowly, or the changes can be detected, we can select/detect periods in which the signal is locally-stationary. We can then compute the spectrum for these periods. Appropriately selecting time periods depends on understanding a fundamental time-frequency uncertainty relationship: The product of the time resolution of the spectral estimate, T and the frequency resolution of the estimate, W , is always greater than 1 (**Fig 3a**). With a time-window of duration T , the lowest resolvable frequency is $1/T$. Deciding how to trade-off time and frequency resolution is complex, as there may be no single best answer. Resolving signals at low frequencies, such as below 10 Hz, requires frequency resolution on the order of 1 Hz. Time resolution is, at best, one second. Conversely, resolving signals that change during behavior, requires time resolution of say 200 ms. Frequency resolution is then 5 Hz or more. Many spectral estimators are available, each with different statistical properties of bias and variance that vary with the desired time-frequency resolution (**Fig 3b,c**). **S1A** discusses spectral estimation with guidelines for best practices.

Interpreting spectral power: In general, field potential power decays in a power-law like way ($1/f^a$), especially for higher frequencies the forward model in a variety of different ways. The $1/f$ shape of the spectrum can generally be sensitive to the overall firing rate of the underlying population or the number of active neurons, with linear offsets in power resulting from an overall change in the firing rate¹⁵. Neuronal spike waveforms can also appear in the LFP, and if the amplitude of spike waveforms in an LFP recording is large. The LFP spectrum at frequencies above 50 Hz can be especially sensitive to changes in firing rate³⁰. LFP power is highly sensitive to the synchrony of the underlying local and afferent-remote populations so the overall shape of the spectrum may reflect a heterogeneous mixture of temporally-structured signals that spanning a range of frequencies. The steepness of the $1/f^a$ curve can also change due to changes in the membrane time-constant. Membrane time constants can depend on the overall level of activation in the circuit⁶², as well as on the activation of NMDA receptors, because these have much slower EPSPs than AMPA receptors^{15,63}. Another contributing factor is noise. Activity at high and very low frequencies often has poor signal to noise ratio, e.g. due to EMG noise (high frequencies) or electrode / cable movement (low frequencies) attenuating observed changes in the underlying signal.

Interpreting oscillations: In addition to the $1/f$ trend, transient, phasic changes in spectral power can be observed to come and go. Detecting a candidate “oscillatory” event depends on setting a threshold. Interpreting these events as changes in a neuronal source is of great importance. In particular, observations of spectral peaks in activity at particular frequencies are often referred to as oscillations. Oscillations are often depicted in the theoretical literature as narrow-band, periodic, sometimes even sinusoidal phenomena. However, theoretical oscillators, like a pendulum generate features at particular frequencies and do not contain multi-component or broadband features. In comparison, physiological neuronal ‘oscillations’ often have a clear broadband or band-limited character^{64,65} and change quickly^{66,67}.

This caveat notwithstanding, we often want to know whether an increase in spectral power indicates an increase in the number of active neurons or an increase in temporally-structured/synchronized neuronal activity. The absence of a spectral peak does not necessarily mean that there is no synchronized activity in the underlying neuronal activity. If there is an underlying oscillatory pattern, a peak in the spike-field coherence (SFC) may be present even without any peak in the LFP spectrum⁶⁸. Thus, if spiking is coherent with a field potential, temporally-structured spiking exists and a local source contributes to the field potential^{39,69} and this source may be said to reflect oscillatory activity. **S1B** presents measures that go beyond second-order correlations to capture correlations between different frequencies and phase-amplitude coupling.

Interpreting cellular sources: Interpreting field potential recordings in terms of cellular sources is limited by the ill-posed nature of the inverse model. Clear interpretations are sometimes available when a local neuronal population generates the currents measured by field potential recordings, such as for the hippocampal theta rhythm⁷⁰. Defining local current generators is called current-source density (CSD) analysis. According to Maxwell’s equations and Ohm’s Law for the extracellular medium, the local current generators are given by a second spatial derivative of the LFP signals in all three directions, called the Laplacian⁶. Therefore, CSD analysis requires invasive recordings with electrode arrays with high-contact density, e.g. 50-200 μm inter-

Investigating large-scale brain dynamics

contact spacing or less, (**Fig 4a**)^{2,5,71}. CSD analysis yields the net volume current density that enters (source) or leaves (sink) the extracellular medium through cell membranes (**Fig 4b**). Consequently, CSD analysis is a more local measure than the LFP and is easier to interpret in terms of neuronal activity.

Traditional CSD analysis assumes that there is no variation of the neuronal activity in the horizontal directions. Thus, CSD is given by the double spatial derivative in the depth direction. This approximation can be problematic as in primary sensory systems, inputs to cortex can be quite focal and significant tangential currents can exist due to tissue curvature (**Fig 4c**), but new CSD estimation methods can account for this. Traditional CSD analysis also assumes that the electrodes adequately sample the currents within the volume. Ideally, three-dimensional sampling of the volume would be preferred to reduce CSD estimation error using the second-spatial derivative method⁶. In general, interpreting current sources in terms of cellular sources, such as whether a sink/source is active (e.g. due to synaptic current) or passive (a return current), depends on additional information about the anatomical connectivity of presynaptic input-generating populations and geometry of postsynaptic populations. **Box 2** and **S1C** present a more detailed discussion.

C2 - Correlation

Networks of neurons generate brain dynamics, and correlated patterns of activity are ubiquitous. There are many ways to capture dependencies between neuronal signals. Here, we will focus on linear measures of coupling, correlations that are well-understood in statistical terms and are in widespread use.

Correlation and coherency: The cross-correlation function measures correlation in the time-domain. In the frequency-domain, we can use the spectral coherence function that is defined as the cross-spectrum between signals normalized by the spectrum of each signal. Coherency is the correlation coefficient that measures the strength of linear association as a complex-valued regression coefficient. The magnitude of the coherence is proportional to the strength of linear association. The phase of the coherence reflects the relative timing but not necessarily a time-delay (see **S2A**).

Like the spectrum and the auto-correlation function, the coherency and the cross-correlation function offer complementary information. An important advantage of the coherency is that it is normalized by power at each frequency. The cross-correlation function is normalized by total variance of each signal which is the integrated power over all frequencies. As a result, the coherency can often detect frequency-localized effects better than the cross-correlation function. The latter is typically dominated by frequencies with more signal power, which are often at lower frequencies.

Estimating coherency depends on similar factors as estimating the spectrum. In general, coherence estimates typically require more DOFs than spectrum estimates. In effect, coherence measures the correlation present in a scatterplot of points, where there is one point for each pair of DOFs. Unlike the spectrum, the distribution of spectral estimators for the coherence is, in general, complicated and depends on the strength of coherence. Several strategies exist to address these issues and statistical tests should be based on appropriate permutation tests^{72,73}.

Two processes that do not interact can appear to be coherent by virtue of being non-stationary⁷⁴. Estimating coherency and cross-correlation in locally stationary time intervals to give time-varying coherence spectra, called coherograms is, therefore, critical. Since coherence estimates with small numbers of DOFs are positively-biased, particular care should be taken when interpreting coherence over short time periods using small numbers of DOF.

Spike-field coherence: Whenever possible, interpreting field potentials in terms of cellular sources is best done directly, using simultaneous measurements of spiking activity. The spectral analysis of spike-field relations largely matches that of field potentials. This is because point processes admit a spectral representation like other time series. As a result, spike-field coherence (SFC) is an important tool for understanding the relationships between LFPs and underlying spiking activity⁵⁸. SFC is the correlation coefficient of a regression, analogous to the coherence between two field potentials⁷⁵, that quantifies how predictable LFP activity is as a linear function of the spike times. SFC specifically measures how neurons tend

Investigating large-scale brain dynamics

to fire spikes at particular phases of LFP activity. The spike-triggered LFP measures spike-field relationships in the time domain and is useful for resolving time-delays between spiking and events in the LFP. Statistical considerations mean the SFC is preferred over the spike-triggered average (see **S2B**).

SFC sensitively reveals the locking of neurons to synchronized synaptic inputs, which also lead to post-synaptic currents and thereby the LFP signal. The broad-band properties of the LFP spectrum itself may mask spectral peaks. However, if there is an underlying oscillatory pattern, a peak in the spike-field coherence (SFC) may be present even without any peak in the LFP spectrum^{76–78}.

Contamination of the LFP by action potentials is a major concern when analyzing phase correlations and SFC. Attenuation of neuronal spike contamination in LFPs can be attempted by removing spikes from the LFP signal⁷⁹. However, residual spike contamination effects may remain. In general, best practice is to use spike and LFP recordings from different electrodes separated by several hundred micrometers.

SNR confound: A critical issue when interpreting correlation and coherence is that both measures are influenced by the relative weight of neuronal signals (signal components) versus other signals (noise components). This leads to an important confound: A change in the signal-to-noise ratio (SNR) will change the measured correlations without a change in the underlying neuronal interaction or functional connectivity. If signal components are more correlated than noise components, e.g. because of a specific interaction between neuronal populations of interest, an increase in signal amplitude or a decrease of noise amplitude can also increase observed correlations. If noise components are more correlated than signal components, e.g. because of field spread from remote neuronal sources to two signals, an increase in noise or a decrease in signal amplitude can increase observed correlations. Correspondingly, changes in noise can also lead to decreases in observed correlations without changes in the underlying neuronal interaction. **S2A** illustrates these effects mathematically.

One strategy to limit concern is to check for changes in signal amplitude that may reflect a potential change in SNR and to only focus on changes in dependencies between signals that either are not paralleled by changes in signal amplitude⁸⁰ or cannot otherwise be accounted for by changes of SNR. Another strategy is to stratify the activity by amplitude. Similarly, the power or SNR of the signal of interest can be binned and correlations can be computed within each bin.

SNR confounds also affect the interpretation of SFC. Enhanced spike-LFP coupling when LFP power increases could reflect greater contribution of spike-locked neuronal sources to the LFP instead of stronger spike-locking of these sources. Differences in mean spiking rates across conditions can also affect the corresponding SFC estimates without requiring true changes in functional connectivity⁸⁰. To mitigate these effects on SFC, spike trains can be decimated to equalize the mean rates across conditions. This involves randomly deleting spike events to match the total number of spikes across conditions. Spike events that are deleted randomly will not alter the underlying spike-field coupling⁸¹. A complementary parametric approach to SFC based on point process Generalized Linear Models (GLMs)⁸² can also be used to separate the contribution of changes in firing rates and changes in phase coupling to LFP oscillations when assessing spike-field relationships across different tasks or conditions^{83,84}. Finally, SFC quantification based on the pairwise phase consistency (PPC) metric avoids spike rate and spike count biases⁸⁵. With this approach, the synchronization between spikes and LFP is determined by first estimating the LFP phase at the time of each individual spike, and then quantifying the similarity of pairs of spike phases.

Volume conduction: Coherence is particularly sensitive to volume conduction and common modes due to distant sources. Volume conduction is of particular concern in the rodent brain. In the rodent, theta and beta frequency range potentials generated by the hippocampal formation can be measured across the neocortical and subcortical neuropil which makes it difficult to perform valid coherence analysis between brain areas^{86,87}.

One effective strategy assumes a forward model in which the impedance of brain tissue is resistive and not capacitive²¹. Field spread due to volume conduction is then effectively instantaneous and does not alter the imaginary component of the cross-spectral density. This yields several measures such as the imaginary part of coherency⁸⁸, phase lag index (PLI)⁸⁹ and weighted phase lag index (WPLI)⁹⁰ (see **S2C**). Best practice is to

Investigating large-scale brain dynamics

analyze activity at each site to assess the presence of volume conduction before interpreting the correlated activity patterns.

Correlations due to volume conduction can survive source reconstruction^{91,92} and re-referencing procedures. The widely-used common-average reference averages all recorded signals and thereby introduces artifacts to which correlations are particularly sensitive. While careful selection of a reference electrode can help suppress shared signals (see **S2D**), care should be taken to assess whether the re-referencing actually removes the common reference. We can also perform spatial filtering to suppress volume conduction by making use of a bipolar reference, computing the CSD or Laplacian and applying source reconstruction / beam-forming techniques described above. The CSD or Laplacian method removes very large-scale, regional or global coherence as well as volume conduction.

C3 – Communication

Given evidence of correlations between two sites in a network, X and Y, we can ask whether the correlations reflect putatively causal interactions with a particular directionality, e.g. X drives Y or $X \rightarrow Y$. Many causal inference procedures exist. Here, we focus on the analysis of Granger Causality (GC) because it is relatively simple and well understood, derives from spectral analysis and is in widespread use.

We should first make clear that, in general, no statistical procedure can accurately recover causal influences if unobserved nodes in the network, or hidden nodes, generate the main interaction effects. This is true for spiking and field potential recordings. Consider a three-node network with common input: If $X \rightarrow Y$ and $X \rightarrow Z$ with a longer delay, unless X is measured, statistical inferences will likely report $Y \rightarrow Z$. Consider another network $X \rightarrow Y \rightarrow Z$. In this case, unless Y is measured, the influence may be incorrectly estimated as $X \rightarrow Z$. Field potentials reflect sub-threshold currents, but the transfer of those currents to spike output can vary from case to case complicating causal inferences. Thus, an influence of field potentials recorded in X on field potentials recorded in Y suggests that spike output of neurons in X influences current flows in neurons in Y, because only spikes travel from X to Y; yet this does not necessarily mean that spike output in Y is influenced. **Box 2** analyzes this topic.

Statistical estimation: Since estimating time delays from correlation/coherence functions can be confounded, additional measures are needed to infer causality and directed influences. Measures based on Granger Causality (GC)^{93,94} assess the direction of influence or causality in terms of temporal prediction among stochastic processes (**Fig 5**). Informally stated, a stochastic process X is said to “Granger-cause” or drive a stochastic process Y if the history of X adds to the prediction of Y beyond what can be predicted based only on the history of Y itself^{95,96}. GC can be statistically assessed in terms of likelihood ratio tests^{97,98} and can be computed both in the time and frequency domains. In addition, if multiple neuronal groups have been recorded simultaneously, the conditional GC can distinguish between: a) direct versus indirect interactions, i.e. $X \rightarrow Z$ vs $X \rightarrow Y \rightarrow Z$, corresponding, for example, to mono vs polysynaptic interactions; or b) between common inputs with different delays vs direct interaction, i.e. between $X \rightarrow Y$ and $X \rightarrow Z$ with a longer delay vs the direct interaction $Y \rightarrow Z$. Software packages to compute GC measures are publicly-available⁹⁸.

Since GC measures typically depend on parametric models, model mismatch is a concern⁹⁹. While GC measures using non-parametric spectral estimates have been proposed¹⁰⁰, a parametric model derived from the non-parametric estimates is still needed. Non-parametric estimates are preferred because they have improved bias and variance properties compared with parametric estimates (see S1). The GC influence measure is also subject to a signal to noise ratio concern because the GC influence assumes there is no measurement noise (see **S3A**). GC estimates also tend to be strongly positively biased by sample size, and re-estimating the GC after shuffling the time series may be needed to control for sample size bias.

Like other spectral analyses, GC analyses can be extended to address slow non-stationarity by using a moving window. Causal inferences depend on model fitting, and non-stationarity can lead to model fitting and interpretation problems for causal inferences. For example, the evoked or event-related potential averaged across trials is subtracted from each trial to ensure stationarity or to separate, assuming a linear additive model, the signal from ongoing background activity. However, if slow across-trial non-stationarities are present

Investigating large-scale brain dynamics

due to changes in neuronal excitability, attention, fatigue, the evoked response can show trial-to-trial changes in amplitude, latency and waveform (see **S1A**). This non-stationarity will affect the interpretation of GC as well as other spectral measures such as power and coherence^{101–105}. Amplitude changes can be effectively suppressed by fitting the evoked response per trial^{106,107}. Latency and waveform changes are more complicated to model, and removal of trial-to-trial variability due to these changes should be performed with caution.

Causal inferences ultimately require empirical validation using interventions. This said, inferences based on interventions also require caution. Interventions often shift the system out of the normal physiological range. Since the intended and achieved manipulations can differ, interventions can also lack validity. Electrical microstimulation preferentially stimulates axons not cell bodies¹⁰⁸. Optogenetic stimulation affects illuminated neurons simultaneously, potentially changing phase delays across layers¹⁰⁹. In general, experiments should use small perturbations to avoid driving the system out of range. Experimental designs should also parametrically vary manipulation conditions instead of simply comparing manipulation and no manipulation conditions to more directly infer the underlying causal mechanisms.

C4 - Coding

Understanding nervous system function depends on the functional forward model - how information is present in activity, what form it takes, and whether and how it can support behavior. Coding implies the presence of correlations between a neuronal signal and external events, such as stimulus onset, and internal events, like the spike of a neuron or the phase of another neuronal signal. Neural decoding quantifies the information present in those correlations.

Information about the task-relevant variables can be decoded from field potential amplitude¹¹⁰, power^{39,42}, coherency magnitude¹¹¹ and phase⁷⁷ and phase-amplitude coupling¹¹². Information can also be decoded using the phase relations between spikes and the surrounding LFP, SFC, as well as by using GLMs that relate spiking probability to the amplitude and phase of LFP oscillations⁸⁴. In the hippocampus, the SFC is strong in the theta range, and its phase carries information about the position of the animal¹¹³. Phase relations between LFPs carry similar information (**Fig 6a**)¹¹⁴. During short-term memory, SFC in the beta band (~ 30 Hz) underlies phase-dependent coding of short-term memory information in prefrontal cortex¹¹⁵. Different memory items are preferentially encoded at different 30 Hz phases (**Fig 6b**). Posterior parietal and frontal neurons that display coherent phase relations, termed coherent neurons, encode information about behaviorally-relevant processes such as decision making and coordination more accurately than neurons that do not fire spikes coherently^{69,116,117}. Information about movement choice is also modulated by LFP phase¹¹⁸ (**Fig 6c**). In visual cortex during visual stimulation, SFC is strong in the gamma band, and its phase depends systematically on stimulus orientation¹¹⁹. Spikes occur earlier in the gamma cycle for stimuli close to the preferred orientation of the spiking neuron than for non-preferred stimuli (**Fig 6d**). In general, phase relations in the gamma frequency band are diverse and change with stimulation and also e.g. with selective attention⁷⁷.

Neuronal decoding analyses explicitly test whether our certainty about the value of a random variable is altered by knowledge of another signal. Modern tools for statistical inference, also called machine learning, are increasingly able to detect subtle patterns in large volumes of data^{49,50,52,120,121}. Over-fitting can inadvertently occur when parameters and other features in the model, often called hyper-parameters or meta-parameters, are selected across the dataset¹²⁰. Cross-validation is the basic technique to address over-estimation of decoding performance due to over-fitting. Problems arise when the statistical model is very flexible and involves making many decisions during model fitting. Strictly speaking, no part of the data available during any stage of fitting should be used to estimate generalization performance. A good practice in decoding analyses is to divide the data into three subsets: a training set to fit a decoding model, a validation set to compare different choices of hyper-parameters or model type, and a test set to assess the decoding performance of the model that has been optimized on the training/validation sets.

Another common issue when estimating neuronal decoding performance is to use features that have been estimated with insufficient statistical DOF. The resulting estimation noise degrades signal coding. In addition, since single-trial measures contain few DOFs, measures of power, correlation and other aspects of brain

Investigating large-scale brain dynamics

dynamics are averaged, typically, across multiple trials. This makes it difficult to measure neuronal coding and how coherence changes trial-by-trial with, for example, reaction times, attention or choice^{116,122,123}. One approach is to perform decoding on the relative phase between signals instead, or derive single-trial pseudo-estimates of coherence through jack-knifing¹²³. The jackknife is more sensitive and accurate than approaches based on sorting-and-binning of trials¹²⁴. Another strategy is to use a classification approach and decode activity on groups of trials with similar single trial performance as-if they were simultaneously-recorded¹¹⁶.

The analysis of neuronal coding in field potentials should not be conflated with the mechanisms by which neurons process information and compute. Field potentials can in some cases influence neuronal information processing due to ephaptic coupling^{125,126}. However, analyses to decode field potentials neither imply nor depend on whether ephaptic coupling comprises the underlying mechanism. The ability to decode information from field potentials can have mechanistic implications.

Phase relations in field potentials may reflect levels of neuronal excitability, may carry behaviorally-relevant information and may influence neuronal communication. Theoretical studies suggest that the phase-leading neuronal activity exerts a stronger influence over the lagging one than vice versa¹²⁷. Experimentally, the phase-leading recording site shows a transfer entropy to the phase-lagging site¹²⁸. Naturally, care is needed when making causal inferences. Phase precession, where the position of the rodent and phase of firing of a place cell correlates with theta LFP activity, is an important link between population coding and oscillatory dynamics. However, information in spiking about spatial position according to theta phase does not necessarily mean the signal has a causal or mechanistic role in the underlying computation. Coding may simply reflect correlation with a latent cause. This concern is a general one and it also applies to inferring a mechanistic role for coding by single neurons where none may be present^{129,130}.

D - Conclusion

We have discussed strategies for analysing and interpreting a range of field potential recording modalities spanning microns to centimeters. The forward model is the central construction. Biophysics, geometry and anatomical connectivity of neuronal populations are the principal ingredients. Several obstacles hinder future progress. One important direction is to simulate virtual experiments using computational models. In these experiments, the “ground truth” is known and the measurement projection from activity space can be defined. Data from virtual experiments can generate model-based benchmarking data. Such benchmarks can test the effects of electrode placement, signal processing, and guide inferences such as directed interactions. The virtual experimental paradigm has been pursued to test blind-source separation algorithms with promising results^{131,132}. Tools to compute field potentials from structured point-neuron networks, such as hybrid LFPy package¹³², can help perform such experiments. Another important direction is to strengthen the forward model by more precisely measuring biological ground truth. For example, genetic tools and microscopy can increasingly reconstruct anatomical information either *in vivo* during an experiment¹³³, or *ex vivo* following the experiment^{134,135}. Inverse models that are sufficiently constrained by ground-truth benchmarking data and empirical observations will permit increasingly rigorous inferences from field potential recordings.

How to validate causal inferences is another recurring theme. Valid causal inferences are powerful because when experimentally controlling a variable, $x(t)$, we can exclude the possibility that correlations between $x(t)$ and an effect, $y(t)$, are explained by a third variable that drives fluctuations in both signals. Simply silencing or lesioning a population of neurons offers a relatively crude picture of causal influences between brain regions. Silencing $x(t)$ may reduce power in another area $y(t)$ but a deeper understanding is masked by the absence of $x(t)$. A more fruitful avenue is to apply relatively small perturbations that allow one to make comparisons with a model fit based on experimental observations. A detailed network-based forward model to design of network control experiments of this kind will be particularly useful for inferring causal mechanisms.

More accurate forward models and analyses informed by inverse models will lead to greater understanding of large-scale brain functions, the design of better electrode arrays and other sensors, and more precise and effective clinical treatments. We hope to have provided a framework that will guide current and future work by researchers who will contribute to these important goals.

Figure Captions

Figure 1 –Field potential recording modalities:

a) EEG and MEG signals are measured non-invasively. EEG involves electrodes ~10 mm in size placed at the scalp across the head. MEG is measured using sensitive sensors (SQUIDS) placed just outside the head¹⁰. ECoG is measured invasively and involves placing electrodes either epidurally on the dura that protects the brain, or sub-durally, directly on the pia at the surface of the brain. ECoG can be recorded in humans in the relatively rare case of epilepsy surgery and is otherwise mainly used in animal models. ECoG electrodes are smaller than EEG electrodes and range in size from one to several millimeters in size¹³⁶.

b) Invasive recordings can also be made at finer spatial scales. Micro-ECoG involves 20-200 μm contacts placed on the pia^{137,138}. Coverage can extend to many square centimeters at sites across the brain. LFP is the most invasive procedure and involves inserting electrodes into the brain. As a result, LFP recordings are made with even smaller size recording contacts than ECoG, extending to microelectrodes and thin-film electrodes that can also record the activity of individual neurons.

Figure 2 – Forward models:

a) Biophysical forward model

The biophysical forward model predicts the amplitude of field potentials generated by populations of neurons depends on dendritic morphology, somato-dendritic location of the incoming synaptic inputs and not axonal morphology of local neurons. **Box 1** discusses other contributions.

i) L4 stellate neurons have restricted and symmetric dendritic arbors (blue) with extended axonal distributions that ramify locally (red). L5 pyramidal neurons have the most extended dendritic arbors and relatively sparse axonal distributions. Adapted from Shepherd et al (2005)¹³⁹. Synaptic input to a population of neurons generates a current dipole that gives rise to the extracellular field potential signal according to the biophysical forward model. **ii)** Populations of L5 pyramidal neurons can receive synaptic input to the apical or basal dendrites and generate large amplitude field potentials in each case. **iii)** Populations of L4 stellate neurons can receive synaptic input near the cell body and generate relatively small amplitude field potentials. Note that this is a simplification. The size of the generated potentials will depend on how displaced the return currents are from the synaptic input currents and spatial distribution of the return currents. The return currents depend on several other factors including the thickness and branching of the dendrites. For example, in some cases about half of the current injected into the apical dendrite of a L5 neuron will return through the soma, and even less in neurons with a thinner apical dendrite¹⁴⁰. Density and strength of synaptic inputs, membrane potential and membrane conductance of the postsynaptic population will also contribute to the magnitude of the generated potentials¹⁴¹.

b) Functional variant of the biophysical forward model.

The functional variant predicts that information can be decoded from field potentials depending on the measurement projection given by the biophysical forward model and the spatiotemporal distribution of source selectivity in the brain.

i) The measurement projection underlying a field potential recording pools information across a volume (shaded). Spatial aspects are illustrated here, but note that the measurement projection also depends on temporal correlations (see **Box 1**). The selectivity of the underlying neuronal sources can be expressed as a directional arrow. **ii)** Field potentials will exhibit strong selectivity if the spatial distribution of the underlying source selectivity is organized so that the measurement projection pools one or several sources with similar selectivity. **iii)** Field potentials will exhibit weak or no selectivity if the spatial distribution of the underlying

source selectivity is disordered and if the measurement projection pools several or many sources. Note that even if the spatial distribution of the underlying source selectivity is organized, weak selectivity can arise if the measurement projection pools over a much larger spatial extent.

Figure 3 – Time series and spectral estimation

a) Fundamentals: **i)** Terms. The sampling rate of a time series, F_s , is inversely related to the sampling interval, dt . The maximum frequency that can be resolved in a time series, called the Nyquist frequency, is determined by the sampling interval and equals $F_s/2$. The bandwidth W is the lowest frequency that can be resolved in a time series, also called the Rayleigh resolution, is inversely related to the duration of the observation window and equals $1/T$. **ii)** Time-frequency plane. The Rayleigh resolution is an example of the time-frequency uncertainty principle which states that the product of the resolution in time T and resolution in frequency $2W$, must be equal or greater than 1, i.e. $2WT \geq 1$. We can increase the time-bandwidth product by increasing the analysis interval T , and hence smoothing more in time, or by increasing bandwidth, W , and smoothing more in frequency. In each case, we effectively assume the properties of neuronal activity are constant within the chosen time and frequency resolution that leads to a smoother, less variable estimate. From Pesaran (2008)⁵⁷.

b) Spectrum of LFP activity in macaque posterior parietal cortex: **i)** Single trial, 500 ms periodogram spectrum estimate. **ii)** Single trial, 500 ms, 10 Hz multitaper spectrum estimate. From Pesaran (2008)⁵⁷.

c) Spectrogram of LFP activity in macaque posterior parietal cortex averaged across 9 trials: Each trial is aligned to the presentation of a spatial cue, which occurs at 0 s. Saccade and reach are made at around 1.2 s. **i)** Multitaper estimate with duration of 500 ms and bandwidth of 10 Hz. **ii)** Multitaper estimate with duration of 200 ms and bandwidth 25 Hz. White rectangle shows the time-frequency resolution of each spectrogram. The color bar shows the spectral power on a log scale in arbitrary units. From Pesaran (2008)⁵⁷.

Figure 4 – Current source density analysis:

a) A cylindrical model in infinite medium. The relationship between current source density (CSD), current flow, and field potential for the simple case of a columnar geometry receiving input at a particular layer. The geometry is a cylinder having a diameter-to-length ratio of 0.5 embedded within an infinite medium of finite conductivity. Due to circular symmetry, only the points on the z-axis along the cylinder are shown. The CSD (I_m ; **i**) gives rise to a current flow (J_z ; **ii**). Current flow establishes the field potentials (V_z ; **iii**). Current density enters the cylinder and returns below. This generates a downward current flow above and below the source/sink and an upward current flow between the source/sink. The resulting field potentials extend beyond the generating current source. Thus, CSD has higher spatial resolution than field potentials. From Nicholson and Freeman (1975)².

b) Laminar patterns of auditory responses in the auditory cortex. **i)** Line plots showing the LFP responses recorded using a linear array multielectrode with 100 μm inter-contact spacing (schematic on left). **ii)** Color plots showing the laminar LFP profile shown in (i). Negative deflections - red. Positive deflections - blue. **iii)** CSD profile estimated using the second spatial-derivative of the field potential profile in (i) and (ii). Red depicts extracellular current sinks associated with net local inward transmembrane current flow. Blue depicts extracellular current sources associated with net local outward trans-membrane current flow. Selected multiunit activity responses are superimposed on the CSD plot. Vertical thin lines indicate stimulus onset. Here, the peak of multiunit activity corresponds to the peak negativity of the LFP and the current sink (CSD) at the response onset in Layer 4. Asterisk indicates a superficial sink that produced the N50 feature in the LFP measured in superficial sites. From Kajikawa and Schroeder (2011)¹⁴².

c) Limitations of traditional CSD estimation method: Simulated one-dimensional recordings for simplified CSD profile in (a). Estimated CSDs for increasing diameter-to-height ratios, as indicated by the number and

Investigating large-scale brain dynamics

inset in the lower left of each panel. Diameter refers to the diameter of the source. All estimates are based on the second spatial-derivative formula of the traditional CSD method. Arbitrary units, negative values to the left and positive to the right. Spurious sources and sinks are inferred for small diameter activity. Estimation error stems from the incorrect assumption of an infinite activity diameter perpendicular to the laminar electrode. To avoid spurious source/sink inferences, a full three-dimensional CSD analysis is needed. Such an analysis can be achieved by making measurements of field potentials in all three spatial directions¹⁴³ and using the second-spatial derivative methods. Alternatively, spurious source and sink inferences can also be avoided by using prior constraints/assumptions as is done in the iCSD¹⁴⁴ and kCSD¹⁴⁵ methods. From Pettersen et al (2012)¹⁴⁶.

Figure 5 – Granger-causal inferences:

a) Schema of multivariate autoregressive models. Each time series is modeled as a linear combination of its own past, the past of the other time series, and an innovation term. These signals are corrupted by measurement noise that can be either correlated or uncorrelated.

b) Cross-correlation function example. An example showing that cross-correlation methods are not appropriate for detecting causal relationships. In this case, the cross-correlation function tells us that future (lagging) but not past (leading) values of $y(t)$ are strongly correlated with $x(t)$. Despite this, the Granger-causal influence from $y(t)$ to $x(t)$ is stronger than the influence from $x(t)$ to $y(t)$.

c) Application of reversed GC testing to LFP data from monkey V1 and V4. Interaction in the gamma band shows full reversal with time series reversal and is therefore robust. This behavior is shown across V1-V4 pairs. In the beta range, GC causality does not reverse with time series reversal, indicating a potential influence of correlated noise.

Figure 6 – Phase-dependent neuronal coding

a) Place information encoded by the phase of hippocampal LFP activity. **i)** Feature-tuned field potentials (FFPs) recorded from the hippocampus during running on a linear track. FFPs uniformly tile the length of a linear track. The spatial extent and spacing of different FFPs is largely homogeneous across the track (middle). FFPs exhibit phase precession with respect to the first principal component of LFP activity (bottom). **ii)** Activation of FFPs during running in a T-maze. Waiting area is enclosed in a red box. Right panels show close-up of activations in waiting area, separated by direction of entry. Asterisks mark activations that are entry-direction selective. Each point represents a time bin where FFP activation exceeds a threshold, its size indicating the magnitude of activation. In (i), place-field hues are assigned based on location of maximal activation; in (ii), hues are assigned to distinguish neighbors. From Agarwal et al (2014)¹¹⁴.

b) Object information in PFC spiking depends on LFP phase. Phase-dependent information during an object working memory task. PFC neuron spiking encodes the identity of two sequentially presented objects during a delay interval. Spikes carry most information about the memorized objects at specific phases of the local 32 Hz LFP. **i)** Optimal encoding of the first presented object is significantly earlier on the falling flank of the 32 Hz cycle. **ii)** Encoding of the second presented object occurs later (Permutation test, $p=0.007$). Error bars denote SEM. Phase dependence induced by stimulus-locked responses was discounted. Adapted from Siegel et al. (2009)¹¹⁵.

c) Choice information in PPC spiking depends on beta and gamma LFP phases. **i)** Average phase-dependent histogram of spike count for the beta frequency range. Coloring of the phase bins in all histograms corresponds to the schematic phase binning shown in the center. The spike-preferred phase (dark green) is depicted as trough in the schematic to capture the tendency of spiking to occur at or near the troughs of LFP activity. The green bin at 0 corresponds to the average spike-preferred phase in the 200 – 1000 ms epoch after target onset when the choice can be made. The radial distance for each phase bin indicates the difference in spike count to random phases. Error bars depict 95% confidence intervals. The radial black line depicts the trigonometric moment of the histogram with its 95% confidence interval indicated at the end of the line. **ii)**

Investigating large-scale brain dynamics

Same graphical depiction as in (i) but for mutual information about choice (choice-MI). Same data as in (i) at each phase bin. Fully colored circles indicate choice-MI significantly different than the average choice-MI across all phase bins (Permutation test, $p < 0.05$). **iii)** Same as (i) for the gamma frequency range. **iv)** Same as (ii) for the gamma frequency range. From Hawellek et al (2016)¹¹⁸.

d) Orientation information in V1 spiking depends on gamma LFP phases. Match between stimulus orientation and neuronal orientation preference determines spike phase in the gamma cycle. Data from one example V1 neuron. **i)** Firing rate as a function of stimulus orientation. **ii)** The black sine wave at the top and the sinusoidal gray shading in the background illustrate the LFP gamma phase. The colored lines show spike densities as a function of phase in the gamma cycle. The colors correspond to the colors used in the firing rate panel on the left. All spike density curves are probability densities, normalized such that the mean value of each curve is 1/2 (bottom left calibration bar applies to all curves, and curves are offset along the y-axis to correspond to i). From Vinck et al (2010)¹¹⁹.

Box 1 - Biophysics of extracellular potentials

Transmembrane currents generate the electrical and magnetic signals recorded from the brain⁷¹. These currents pass through the membranes of neurons and glial cells due to active mechanisms, as well as passive mechanisms involving capacitive coupling between conductive elements. *Volume-conductor theory* explains how signals propagate from their sources in cells through brain tissue to the electrode^{6,10}. Volume-conductor theory provides the biophysical basis for recorded extracellular signals. For example, it allows us to generate extracellular signals from transmembrane currents by simulating the dynamics of biophysically-detailed neuron or neuronal network models using biophysically-detailed multi-compartment neuron models based on cable theory (**Fig Box 1a**)^{5,147–149}. The *cable theory* of neurons further implies that the sum of the transmembrane ionic and capacitive currents across the entire cellular surface must be zero¹⁵⁰. As a result, currents enter the cell and are balanced by currents that leave at other locations. Note though that even in the absence of transmembrane current there may be extracellular potentials set up by *diffusion* of ions¹⁵¹.

Volume-conductor theory and cable theory allow us, in general terms, to calculate single-neuron currents from morphologically and biophysically-detailed neuron models (**Fig Box 1a**)¹⁴⁰. We can then determine how much each neuron contributes to the relevant field potential signal, i.e., LFP¹² EEG and MEG^{6,10}. According to the theory, field potentials summate activity from all electrically-active membranes and transmembrane current generators in space and time, from axon terminals to soma, from action potentials to very slow conductances. The precise contribution of a given cell depends on geometric factors such as cell morphology and the anatomical distribution of active conductances and synaptic inputs to the cell²⁶. For recordings far from the neuronal source, like M/EEG, we can represent the current sources using the *dipole approximation*. The dipole approximation summarizes the summed contribution of all micro-current sources within a tissue volume by an effective dipole moment per unit volume⁶. Consequently, the forward model depends on the relative orientation and separation of the underlying neuronal sources.

While the strength of the effective dipole moment generally increases with the strength of synaptic micro-sources, geometric factors often dominate. If the micro-sources are symmetrically distributed in space, the strength of the resulting meso-source will tend to be small because the dipoles will cancel each other. This is true even when there are relatively large micro-sources and is known as a “closed field”¹⁵². In contrast, if the positive and negative sources are spatially separated, relatively large meso-sources can form even if the underlying micro-source magnitudes are relatively small. This is known as an “open field”. Pyramidal neurons, the most numerous neurons in the hippocampus and cortex, have extended dendritic arbors and open fields.

While many sources are known to contribute to the field potential, in general, the measurement projection made by a given field potential recording preferentially weights post-synaptic potentials in neurons and brain systems with an open-field geometry and spatially organized synaptic inputs. Synaptic inputs onto either the basal or apical dendrites that are correlated in time will set up the largest extracellular potentials (**Fig Box 1b,c**). If synaptic inputs are evenly distributed, the resulting transmembrane currents will tend to cancel, and the generated extracellular potential will be small^{153,154}. When receiving spatially uniform synaptic input, stellate, basket cells and other neurons with symmetric dendritic arbors form closed fields and contribute relatively little. Myelinated axonal fibers, non-myelinated axon compartments, and presynaptic terminals are also expected to contribute relatively little to LFPs as they have small membrane areas¹⁴⁸. If synaptic inputs are spatially separated, significant current dipoles will form¹⁵³. For example, stellate cells can function as open-field generators when they receive asymmetric synaptic input^{140,155}.

Summation effects extend beyond the individual neuron. Small amplitude current sources that are arranged into particular spatial configurations across neurons, a sheet, for example, can give rise to large-amplitude signals. Interference by return currents in other neurons can also cancel input-related active synaptic currents and reduce signal amplitude. Conversely, constructive alignment of opposite polarities of coherently active dipoles amplifies the juxtaposed pole. In rodent, this can be seen in dentate gyrus with its opposing granular layer blades¹⁵⁶, and likely in other cases, such as bilateral dipoles of the medial cortical areas.

Figure Box 1 – Extracellular potential biophysics:

a) Application of cable theory in a multi-compartmental model. In cable theory, we simplify the three-dimensional complexity of the long, thin dendrites and axons to a one-dimensional core conductor along the long axis. This is the axis along which the membrane potential will vary the most. In this example, an apical dendritic branch, assumed purely passive with only capacitive and leak membrane currents, is divided into a set of compartments indexed by n . The circuit diagram shows the equivalent electric circuit of the compartment. The net transmembrane current $I_n(t)$, is, in this case, the sum over the capacitive and leak membrane currents in compartment n . $I_n(t)$ is then used in forward-modeling schemes, such as is implemented in LFPy¹², to calculate extracellular potentials. Two elements in the equivalent electric circuit represent intracellular resistive currents between compartment n and the neighboring compartments $n+1$ and $n-1$. Other elements represent currents due to capacitive properties of the cell membrane, and various other membrane processes such as passive and active intrinsic ion channels and synaptic inputs. If we assume point current sources, the extracellular potential $\varphi(\mathbf{r}, t)$ recorded at position \mathbf{r} due to each of the transmembrane currents $I_0(t)$ at position \mathbf{r}_0 is given by $\varphi(\mathbf{r}, t) = I_0(t)/4\pi\sigma|\mathbf{r} - \mathbf{r}_0|$. σ is the extracellular conductivity, assumed to be real, scalar and homogeneous. Note that the simplest model producing an extracellular potential is a two-compartment model where a transmembrane current entering the neuron at one compartment leaves at the other compartment and forming a current dipole. From Linden et al (2014)¹².

b) Field potential amplitude depends on neuronal morphology. i) Simulations illustrating the dependence of population LFP on the “pyramidalness” of the neurons, i.e., on the distance between cylinders containing the basal and apical dendrites (apical cylinder marked with transparent blue). When the two cylinders are completely superimposed (left panel), the structure corresponds to a stellate cell. When the two cylinders are positioned immediately on top of each other (center panel), the morphology roughly corresponds to a layer 2/3 pyramidal cell. When the boundaries of the two cylinders are separated by 250 μm (right panel), the cell morphology resembles a layer 5 pyramidal cell. In all cases GABA synapses were distributed only on dendrites in the lower cylinder, while AMPA synapses were distributed over the entire dendritic tree (cf. “Reference” distribution in panel **c**). **ii)** Average absolute amplitude (standard deviation) of LFP fluctuations as a function of distance between cylinders. The LFP value corresponds to the LFP amplitude averaged across depths along the axis of the cylinders. From Mazzoni et al (2015)¹⁴⁹.

c) Field potential amplitude depends on the distribution of synaptic inputs. i) Simulations for different synaptic distributions. **Left:** Homogeneous - both AMPA and GABA synapse distributed over the entire surface of the cell. **Center:** Reference - GABA synapses distributed only in the lower cylinder, with AMPA synapses distributed over the entire cell. **Right:** Separate - GABA synapses distributed only in the lower cylinder and AMPA synapses only in the upper cylinder. **ii)** Average LFP absolute amplitude versus dipole moment (standard deviation over time) for the different synaptic distributions (Homogeneous-black; Reference-red; Separated-green). From Mazzoni et al (2015)¹⁴⁹.

Box 2 – Local and remote sources

A basic question is, “How much do field potentials reflect local population activity as opposed to the activity of remote sources?”. Setting aside for a moment the issue of volume conduction, (see above and S2C), the currents that contribute substantially to the LFP are post-synaptic potentials in nearby neurons (see Box 1). Consistent with this, the LFP correlates closely with nearby intracellular membrane potential recordings^{157,158}.

What remains unstated, however, is whether the synaptic currents generating the LFP originate from local or remote neuronal firing. Post-synaptic potentials that generate local current dipoles will result from the firing of nearby neurons forming local, recurrent connections as well as the firing of remote neurons with afferent inputs into a region. Distinguishing afferent inputs and local activity in general may not be possible. In most cases a mixture of the two should be assumed to contribute to the LFP signal. In addition to recording neurons locally and analyzing SFC, we also need to analyze the post-synaptic consequences of the inputs and the location and properties of our recording contacts. Importantly, both IPSCs and EPSCs can contribute to LFP signals^{4,159}. Perisomatic inhibition can set up a dipole in pyramidal cells and the contribution of GABAergic interneurons to the LFP may be boosted by their high synchrony and divergent projections onto pyramidal cells causing correlated IPSCs in many pyramidal cells¹⁵⁴. Distinct classes of GABAergic populations project specifically to soma or dendritic regions and so generate different current dipoles.

Source-sink locations contain important information about whether activity is mostly locally generated. For example, cortical L2/3 source-sink pairs argue against a contribution of afferents to granular (e.g. thalamic) and infragranular layers. In contrast, L4 source-sink pairs tend to indicate input through a feed-forward projection from the thalamus^{160–162}, though contribution from the local recurrent afferents cannot be excluded. Another strategy to resolve local from distal contributions is to record the major afferent systems together with the local activity¹⁶¹ or to perturb local spiking activity using pharmacological and optogenetic manipulations.

The extent to which afferents and local recurrent activity contribute to field potentials may depend on the brain area, the species, the state of the animal (e.g. wakefulness vs. anesthesia) and the behavioral task. In the cat and the primate, there is abundant evidence showing that visually induced gamma-band synchronization arises from recurrent activity within layer 2/3 and layer 4B^{161,163,164} (**Fig Box 2a**). However, activity in frequency bands below ~30 Hz, including alpha and beta, displays long-range SFC and may not be locally-generated.

Interpreting local/remote attributions is difficult when area X sends strong feed-forward synaptic connections to area Y and when area Y has local recurrent connectivity. Under both conditions, LFPs in area Y can reflect the impact of synaptic input from area X. Area X input may or may not drive spiking in Y due to the effect of frequency-dependent dendritic filtering⁴. The presence of signals from area X on LFPs in area Y only depends on area X having strong anatomical projections to area Y and synchronized neuronal firing in X projecting to Y. The effect of such afferent inputs can be seen clearly in how thalamic spiking causes a response in layer-4/6 in rabbit¹⁶⁵ (**Fig Box 2b**) and cat¹⁶⁶, and how the field response in a single whisker column layer-4 reflects synchronization in the matching thalamic barrelloid^{160,167} (**Fig Box 2c**).

Anatomical projections with highly correlated signals preferentially contribute to field potential recordings¹⁵⁴. For example, hippocampal CA3 neuron gamma frequency output causes postsynaptic currents in the CA1 radiatum, a projection region of CA3^{168,169}. Thus, the field potential in area Y is influenced by both, the local activity in Y as well as the delayed copy of activity in area X. The presence of field coherence between X and Y is not a sufficient evidence for neural population coupling. Further evidence that neurons in area Y change their firing in response to input from X is needed. This caveat also applies to Granger Causal influences. Conversely, absence of field coherence between X and Y is not a sufficient evidence for the lack of neuronal population coupling. The forward model given connections between X and Y may predict a small contribution of X inputs to the LFP in Y, and so generate little coherence.

In general, therefore, simple rules cannot be provided. The contribution of local/remote depends on the area, species and the spectral band. CSD analysis cannot resolve the issue, as it merely distinguishes between volume conducted activity and local dipoles. SFC offers important constraints. The absence of local SFC in a particular frequency band can be used as evidence against local origin. The presence of SFC within

Investigating large-scale brain dynamics

area Y is evidence in support of but not proof of $X \rightarrow Y$ communication. Recordings from the anatomically-projecting afferent systems along with manipulations of local cell activity are needed to provide direct confirmation.

Figure Box 2 – Local and remote sources:

a) Local activity in cat visual cortex. **i)** Moving stimuli generate coherent gamma oscillations in cat visual area 17. **ii)** The same stimuli generate higher-frequency oscillations in LGN. **iii)** Visual cortex and LGN oscillations are not coherent with each other. From Castelo-Branco et al (1998)¹⁷⁰.

b) Gamma oscillations in developing rodent cortex due to thalamic input. **i)** Experimental setup for simultaneous recordings of single-whisker evoked responses in the corresponding VPM barreloid and cortical barrel column in the P6 rat. **ii)** Responses averaged across 100 whisker deflections in the matching barreloid of VPM thalamus (VPM MUA PSTH (black) and granular layer of the barreloid of the cortex (evoked LFP (red))). Cortical LFP gamma activity is present at a stage of development during which cortical circuits do not generate gamma activity. Thalamic multiunit activity preceding cortical gamma implies that cortical LFP is due to incoming thalamic spiking and not local cortical spiking. From Minlebaev et al (2011)¹⁶⁰.

c) LFP activity in rabbit cortex due to thalamic input. **i)** Experimental setup in which two thalamic electrodes recorded from neurons in neighboring VB barreloids (labeled a and b), and cortical electrodes recorded spike-triggered averages in the topographically aligned (and neighboring) S1 barrels (labeled A and B). Spike-triggered averages were generated by two neurons in barreloid a (a1 and a2) and by 1 neuron in barreloid b. **ii)** Spike-triggered averages elicited in barrel A by barreloid neurons a1, a2, and b are shown at bottom left. Spike-triggered averages elicited in barrel B by the same 3 VB neurons is shown at bottom right. From Swadlow and Gusev (2000)¹⁶⁷.

References

1. Leung, L.-W. S. Field Potentials in the Central Nervous System Recording, Analysis, and Modeling. in *Neuromethods, Vol 15: Neurophysiological Techniques: Applications to Neural Systems* (eds. Boulton, A., Baker, G. & Vanderwolf, C.) 277–312 (The Humana Press, Inc, 1990).
2. Nicholson, C. & Freeman, J. A. Theory of current source-density analysis and determination of conductivity tensor for anuran cerebellum. *J Neurophysiol* **38**, 356–368 (1975).
3. Gratiy, S. L. *et al.* From Maxwell's equations to the theory of current-source density analysis. *Eur J Neurosci* **45**, 1013–1023 (2017).
4. Buzsáki, G., Anastassiou, C. A. & Koch, C. The origin of extracellular fields and currents--EEG, ECoG, LFP and spikes. *Nat Rev Neurosci* **13**, 407–20 (2012).
5. Einevoll, G. T., Kayser, C., Logothetis, N. K. & Panzeri, S. Modelling and analysis of local field potentials for studying the function of cortical circuits. *Nat Rev Neurosci* **14**, 770–85 (2013).
6. Nunez, P. & Srinivasan, R. *Electric fields in the Brain. The Neurophysics of EEG.* (Oxford Univ Press, 2006).
7. Hämäläinen, M. S. Magnetoencephalography: a tool for functional brain imaging. *Brain Topogr* **5**, 95–102 (1992).
8. Fries, P. A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. *Trends Cogn Sci* **9**, 474–80 (2005).
9. Siegel, M., Donner, T. H. & Engel, A. K. Spectral fingerprints of large-scale neuronal interactions. *Nat. Rev. Neurosci.* **13**, 121–34 (2012).
10. Hämäläinen, M., Hari, R., Ilmoniemi, R. J., Knuutila, J. & Lounasmaa, O. V. Magnetoencephalography—theory, instrumentation, and applications to noninvasive studies of the working human brain. *Rev Mod Phys* **65**, 413–497 (1993).
11. Nunez, P. L. Neocortical dynamics of macroscopic-scale EEG measurements. *IEEE Eng. Med. Biol. Mag.* **17**, 110–7 (1998).
12. Lindén, H. *et al.* LFPy: a tool for biophysical simulation of extracellular potentials generated by detailed model neurons. *Front Neuroinform* **7**, 41 (2013).
13. Gilja, V. & Moore, T. Electrical signals propagate unbiased in cortex. *Neuron* **55**, 684–6 (2007).
14. Bédard, C., Kröger, H. & Destexhe, A. Model of low-pass filtering of local field potentials in brain tissue. *Phys. Rev. E* **73**, 051911 (2006).
15. Miller, K. J., Sorensen, L. B., Ojemann, J. G. & den Nijs, M. Power-Law Scaling in the Brain Surface Electric Potential. *PLoS Comput. Biol.* **5**, e1000609 (2009).
16. Pettersen, K. H., Lindén, H., Tetzlaff, T. & Einevoll, G. T. Power Laws from Linear Neuronal Cable Theory: Power Spectral Densities of the Soma Potential, Soma Membrane Current and Single-Neuron Contribution to the EEG. *PLoS Comp Biol* **10**, e1003928 (2014).
17. Ranck, J. Specific impedance of rabbit cerebral cortex. *Exp Neurol* **7**, 144–52 (1963).
18. Pfurtscheller, G. & Cooper, R. Frequency dependence of the transmission of the EEG from cortex to scalp. *Electroencephalogr. Clin. Neurophysiol.* **38**, 93–6 (1975).
19. Gabriel, S., Lau, R. W. & Gabriel, C. The dielectric properties of biological tissues: II. Measurements in the frequency range 10 Hz to 20 GHz. *Phys. Med. Biol.* **41**, 2251–69 (1996).
20. Bédard, C. & Destexhe, A. Macroscopic models of local field potentials and the apparent 1/f noise in brain activity. *Biophys J* **96**, 2589–603 (2009).
21. Logothetis, N. K., Kayser, C. & Oeltermann, A. In vivo measurement of cortical impedance spectrum in monkeys: implications for signal propagation. *Neuron* **55**, 809–823 (2007).

Investigating large-scale brain dynamics

22. Wagner, T. *et al.* Impact of brain tissue filtering on neurostimulation fields: a modeling study. *Neuroimage* **85 Pt 3**, 1048–57 (2014).
23. Dowrick, T., Blochet, C. & Holder, D. In vivo bioimpedance measurement of healthy and ischaemic rat brain: implications for stroke imaging using electrical impedance tomography. *Physiol Meas* **36**, 1273–82 (2015).
24. Elbohouty, M., Wilson, M. T., Voss, L. J., Steyn-Ross, D. A. & Hunt, L. A. In vitro electrical conductivity of seizing and non-seizing mouse brain slices at 10 kHz. *Phys Med Biol* **58**, 3599–613 (2013).
25. Miceli, S., Ness, T. V., Einevoll, G. T. & Schubert, D. Impedance Spectrum in Cortical Tissue: Implications for Propagation of LFP Signals on the Microscopic Level. *eNeuro* **4**, ENEURO.0291-16.2016 (2017).
26. Reimann, M. W. *et al.* A biophysically detailed model of neocortical local field potentials predicts the critical role of active membrane currents. *Neuron* **79**, 375–90 (2013).
27. Schomburg, E. W., Anastassiou, C. A., Buzsáki, G. & Koch, C. The spiking component of oscillatory extracellular potentials in the rat hippocampus. *J. Neurosci.* **32**, 11798–811 (2012).
28. Scheffer-Teixeira, R., Belchior, H., Leão, R. N., Ribeiro, S. & Tort, A. B. L. On high-frequency field oscillations (>100 Hz) and the spectral leakage of spiking activity. *J. Neurosci.* **33**, 1535–9 (2013).
29. Ray, S. & Maunsell, J. H. R. Different origins of gamma rhythm and high-gamma activity in macaque visual cortex. *PLoS Biol.* **9**, e1000610 (2011).
30. Waldert, S., Lemon, R. N. & Kraskov, A. Influence of spiking activity on cortical local field potentials. *J Physiol* **591**, 5291–303 (2013).
31. Cuffin, B. N. *et al.* Tests of EEG localization accuracy using implanted sources in the human brain. *Ann. Neurol.* **29**, 132–138 (1991).
32. Van Veen, B. D., van Drongelen, W., Yuchtman, M. & Suzuki, A. Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Trans Biomed Eng* **44**, 867–80 (1997).
33. Hipp, J. F., Engel, A. K. & Siegel, M. Oscillatory Synchronization in Large-Scale Cortical Networks Predicts Perception. *Neuron* **69**, 387–396 (2011).
34. Hari, R. & Forss, N. Magnetoencephalography in the study of human somatosensory cortical processing. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **354**, 1145–54 (1999).
35. Hipp, J. F. & Siegel, M. Dissociating neuronal gamma-band activity from cranial and ocular muscle activity in EEG. *Front Hum Neurosci* **7**, 338 (2013).
36. Krings, T., Chiappa, K. H., Cuffin, B. N., Buchbinder, B. R. & Cosgrove, G. R. Accuracy of electroencephalographic dipole localization of epileptiform activities associated with focal brain lesions. *Ann. Neurol.* **44**, 76–86 (1998).
37. Riera, J. J. *et al.* Pitfalls in the dipolar model for the neocortical EEG sources. *J Neurophysiol* **108**, 956–75 (2012).
38. Barth, D. S. Empirical comparison of the MEG and EEG: Animal models of the direct cortical response and epileptiform activity in neocortex. *Brain Topogr* **4**, 85–93 (1991).
39. Pesaran, B., Pezaris, J. S., Sahani, M., Mitra, P. P. & Andersen, R. A. Temporal structure in neuronal activity during working memory in macaque parietal cortex. *Nat Neurosci* **5**, 805–811 (2002).
40. Bansal, A. K., Truccolo, W., Vargas-Irwin, C. E. & Donoghue, J. P. Decoding 3-D reach and grasp from hybrid signals in motor and premotor cortices: spikes, multiunit activity and local field potentials. *J Neurophysiol* (2011). doi:10.1152/jn.00781.2011
41. Katzner, S. *et al.* Local origin of field potentials in visual cortex. *Neuron* **61**, 35–41 (2009).
42. Siegel, M. & König, P. A functional gamma-band defined by stimulus-dependent synchronization in area 18 of awake behaving cats. *J Neurosci* **23**, 4251–4260 (2003).

Investigating large-scale brain dynamics

43. Boes, A. D. *et al.* Network localization of neurological symptoms from focal brain lesions. *Brain* **138**, 3061–3075 (2015).
44. Liu, J. & Newsome, W. T. Local Field Potential in Cortical Area MT: Stimulus Tuning and Behavioral Correlations. *J. Neurosci.* **26**, 7779–7790 (2006).
45. Berens, P., Keliris, G. A., Ecker, A. S., Logothetis, N. K. & Tolias, A. S. Feature selectivity of the gamma-band of the local field potential in primate primary visual cortex. *Front. Neurosci.* **2**, 199–207 (2008).
46. Pistohl, T., Schulze-Bonhage, A., Aertsen, A., Mehring, C. & Ball, T. Decoding natural grasp types from human ECoG. *Neuroimage* (2011). doi:10.1016/j.neuroimage.2011.06.084
47. Cogan, G. B. *et al.* Sensory-motor transformations for speech occur bilaterally. *Nature* **507**, 94–8 (2014).
48. Pesaran, B., Musallam, S. & Andersen, R. A. Cognitive neural prosthetics. *Curr. Biol. CB* **16**, R77-80 (2006).
49. Samaha, J., Sprague, T. C. & Postle, B. R. Decoding and Reconstructing the Focus of Spatial Attention from the Topography of Alpha-band Oscillations. *J Cogn Neurosci* **28**, 1090–1097 (2016).
50. Cichy, R. M., Pantazis, D. & Oliva, A. Resolving human object recognition in space and time. *Nat Neurosci* **17**, 455–62 (2014).
51. Carlson, T. *et al.* Representational dynamics of object vision: The first 1000 ms. *J Vis* **13**, 1–1 (2013).
52. Myers, N. E. *et al.* Testing sensory evidence against mnemonic templates. *Elife* **4**, (2015).
53. Srinivasan, R., Nunez, P. L. & Silberstein, R. B. Spatial filtering and neocortical dynamics: estimates of EEG coherence. *IEEE Trans Biomed Eng* **45**, 814–26 (1998).
54. Rickert, J. *et al.* Encoding of movement direction in different frequency ranges of motor cortical local field potentials. *J. Neurosci.* **25**, 8815–24 (2005).
55. Gunduz, A. *et al.* Decoding covert spatial attention using electrocorticographic (ECoG) signals in humans. *Neuroimage* **60**, (2012).
56. Brillinger, D. R. *Time Series*. (Society for Industrial and Applied Mathematics, 2001). doi:10.1137/1.9780898719246
57. Pesaran, B. Spectral analysis for neural signals. in *Neural Signal Processing: Quantitative Analysis of Neural Activity* (ed. Mitra, P. P.) 1–13 (Society for Neuroscience, 2008).
58. Halliday, D. M. *et al.* A framework for the analysis of mixed time series/point process data--theory and application to the study of physiological tremor, single motor unit discharges and electromyograms. *Prog Biophys Mol Biol* **64**, 237–278 (1995).
59. Aru, J. *et al.* Untangling cross-frequency coupling in neuroscience. *Curr Opin Neurobiol* **31**, 51–61 (2015).
60. Steriade, M., Timofeev, I. & Grenier, F. Natural waking and sleep states: a view from inside neocortical neurons. *J Neurophysiol* **85**, 1969–85 (2001).
61. McGinley, M. J. *et al.* Waking State: Rapid Variations Modulate Neural and Behavioral Responses. *Neuron* **87**, 1143–61 (2015).
62. Destexhe, A., Rudolph, M. & Paré, D. The high-conductance state of neocortical neurons in vivo. *Nat. Rev. Neurosci.* **4**, 739–751 (2003).
63. van Wingerden, M. *et al.* NMDA Receptors Control Cue-Outcome Selectivity and Plasticity of Orbitofrontal Firing Patterns during Associative Stimulus-Reward Learning. *Neuron* **76**, 813–825 (2012).
64. Burns, S. P., Xing, D., Shelley, M. J. & Shapley, R. M. Searching for autocorrelation in the cortical network with a time-frequency analysis of the local field potential. *J Neurosci* **30**, 4033–47 (2010).
65. Burns, S. P., Xing, D. & Shapley, R. M. Is gamma-band activity in the local field potential of V1 cortex a 'clock' or filtered noise? *J Neurosci* **31**, 9658–64 (2011).

Investigating large-scale brain dynamics

66. Rule, M. E., Vargas-Irwin, C. E., Donoghue, J. P. & Truccolo, W. Dissociation between sustained single-neuron spiking and transient β -LFP oscillations in primate motor cortex. *J. Neurophysiol.* **117**, 1524–1543 (2017).
67. Feingold, J., Gibson, D. J., DePasquale, B. & Graybiel, A. M. Bursts of beta oscillation differentiate postperformance activity in the striatum and motor cortex of monkeys performing movement tasks. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 13687–92 (2015).
68. Zeitler, M., Fries, P. & Gielen, S. Assessing neuronal coherence with single-unit, multi-unit, and local field potentials. *Neural Comput.* **18**, 2256–81 (2006).
69. Wong, Y. T., Fabiszak, M. M., Novikov, Y., Daw, N. D. & Pesaran, B. Coherent neuronal ensembles are rapidly recruited when making a look-reach decision. *Nat. Neurosci.* **19**, 327–34 (2016).
70. Ylinen, A. *et al.* Intracellular correlates of hippocampal theta rhythm in identified pyramidal cells, granule cells, and basket cells. *Hippocampus* **5**, 78–90 (1995).
71. Mitzdorf, U. Current source-density method and application in cat cerebral cortex: investigation of evoked potentials and EEG phenomena. *Physiol. Rev.* **65**, 37–100 (1985).
72. Maris, E., Schoffelen, J.-M. & Fries, P. Nonparametric statistical testing of coherence differences. *J Neurosci Methods* **163**, 161–75 (2007).
73. Vinck, M., van Wingerden, M., Womelsdorf, T., Fries, P. & Pennartz, C. M. A. The pairwise phase consistency: a bias-free measure of rhythmic neuronal synchronization. *Neuroimage* **51**, 112–22 (2010).
74. Granger, C. W. J. & Newbold, P. Spurious regressions in econometrics. *J Econom.* **2**, 111–120 (1974).
75. Jarvis, M. R. & Mitra, P. P. Sampling properties of the spectrum and coherency of sequences of action potentials. *Neural Comput.* **13**, 717–49 (2001).
76. Vinck, M., Womelsdorf, T., Buffalo, E. A., Desimone, R. & Fries, P. Attentional modulation of cell-class-specific gamma-band synchronization in awake monkey area v4. *Neuron* **80**, 1077–89 (2013).
77. Maris, E., Womelsdorf, T., Desimone, R. & Fries, P. Rhythmic neuronal synchronization in visual cortex entails spatial phase relation diversity that is modulated by stimulation and attention. *Neuroimage* **74**, 99–116 (2013).
78. Brunet, N. M. *et al.* Stimulus repetition modulates gamma-band synchronization in primate visual cortex. *Proc Natl Acad Sci USA* **111**, 3626–31 (2014).
79. Zanos, T. P., Mineault, P. J. & Pack, C. C. Removal of spurious correlations between spikes and local field potentials. *J Neurophysiol* (2010). doi:10.1152/jn.00642.2010
80. Lepage, K. Q., Kramer, M. A. & Eden, U. T. The dependence of spike field coherence on expected intensity. *Neural Comput* **23**, 2209–41 (2011).
81. Ogata, Y. On Lewis' simulation method for point processes. *IEEE Trans. Inf. Theory* **27**, 23–31 (1981).
82. Truccolo, W., Eden, U. T., Fellows, M. R., Donoghue, J. P. & Brown, E. N. A point process framework for relating neural spiking activity to spiking history, neural ensemble, and extrinsic covariate effects. *J Neurophysiol* **93**, 1074–1089 (2005).
83. Lepage, K. Q. *et al.* A procedure for testing across-condition rhythmic spike-field association change. *J Neurosci Meth* **213**, 43–62 (2013).
84. Rule, M. E., Vargas-Irwin, C., Donoghue, J. P. & Truccolo, W. Contribution of LFP dynamics to single-neuron spiking variability in motor cortex during movement execution. *Front Sys Neurosci* **9**, 89 (2015).
85. Vinck, M., Battaglia, F. P., Womelsdorf, T. & Pennartz, C. Improved measures of phase-coupling between spikes and the Local Field Potential. *J Comp Neurosci* **33**, 53–75 (2012).
86. Sirota, A. *et al.* Entrainment of neocortical neurons and gamma oscillations by the hippocampal theta rhythm. *Neuron* **60**, 683–97 (2008).
87. Vinck, M., Batista-Brito, R., Knoblich, U. & Cardin, J. A. Arousal and locomotion make distinct

- contributions to cortical activity patterns and visual encoding. *Neuron* **86**, 740–54 (2015).
88. Nolte, G. *et al.* Identifying true brain interaction from EEG data using the imaginary part of coherency. *Clin Neurophysiol* **115**, 2292–307 (2004).
 89. Stam, C. J., Nolte, G. & Daffertshofer, A. Phase lag index: assessment of functional connectivity from multi channel EEG and MEG with diminished bias from common sources. *Hum Brain Mapp* **28**, 1178–93 (2007).
 90. Vinck, M., Oostenveld, R., van Wingerden, M., Battaglia, F. & Pennartz, C. M. A. An improved index of phase-synchronization for electrophysiological data in the presence of volume-conduction, noise and sample-size bias. *Neuroimage* **55**, 1548–65 (2011).
 91. Schoffelen, J.-M. & Gross, J. Source connectivity analysis with MEG and EEG. *Hum Brain Mapp* **30**, 1857–65 (2009).
 92. Schoffelen, J.-M., Oostenveld, R. & Fries, P. Imaging the human motor system's beta-band synchronization during isometric contraction. *Neuroimage* **41**, 437–447 (2008).
 93. Kamiński, M., Ding, M., Truccolo, W. A. & Bressler, S. L. Evaluating causal relations in neural systems: granger causality, directed transfer function and statistical assessment of significance. *Biol Cybern* **85**, 145–57 (2001).
 94. Brovelli, A. *et al.* Beta oscillations in a large-scale sensorimotor cortical network: directional influences revealed by Granger causality. *Proc Natl Acad Sci USA* **101**, 9849–54 (2004).
 95. Granger, C. W. J. Investigating Causal Relations by Econometric Models and Cross-spectral Methods. *Econometrica* **37**, 424–438 (1969).
 96. Wiener, N. *The Theory of Prediction. Modern Mathematics for Engineers* **58**, (McGraw-Hill, 1956).
 97. Geweke, J. Measurement of Linear Dependence and Feedback between Multiple Time Series. *J. Am. Stat. Assoc.* (2012).
 98. Barnett, L. & Seth, A. K. The MVGC multivariate Granger causality toolbox: a new approach to Granger-causal inference. *J Neurosci Meth* **223**, 50–68 (2014).
 99. Friston, K. J. *et al.* Granger causality revisited. *Neuroimage* **101**, 796–808 (2014).
 100. Dhamala, M., Rangarajan, G. & Ding, M. Analyzing information flow in brain networks with nonparametric Granger causality. *Neuroimage* **41**, 354–62 (2008).
 101. Banerjee, A., Dean, H. L. & Pesaran, B. Parametric models to relate spike train and LFP dynamics with neural information processing. *Front Comp Neurosci* **6**, 51 (2012).
 102. Truccolo, W. A., Ding, M., Knuth, K. H., Nakamura, R. & Bressler, S. L. Trial-to-trial variability of cortical evoked responses: implications for the analysis of functional connectivity. *Clin Neurophysiol* **113**, 206–226 (2002).
 103. Truccolo, W. *et al.* Estimation of single-trial multicomponent ERPs: differentially variable component analysis (dVCA). *Biol Cybern* **89**, 426–38 (2003).
 104. Knuth, K. H. *et al.* Differentially variable component analysis: Identifying multiple evoked components using trial-to-trial variability. *J Neurophysiol* **95**, 3257–76 (2006).
 105. Wang, X., Chen, Y. & Ding, M. Estimating Granger causality after stimulus onset: a cautionary note. *Neuroimage* **41**, 767–76 (2008).
 106. Xu, L. *et al.* ASEO: A method for the simultaneous estimation of single-trial event-related potentials and ongoing brain activities. *IEEE Trans. Biomed. Eng.* **56**, 111–121 (2009).
 107. Banerjee, A., Dean, H. L. & Pesaran, B. Parametric models to relate spike train and LFP dynamics with neural information processing. *Front. Comput. Neurosci.* **6**, (2012).
 108. McIntyre, C. C. & Grill, W. M. Selective microstimulation of central nervous system neurons. *Ann Biomed Eng* **28**, 219–33 (2000).

Investigating large-scale brain dynamics

109. Buzsáki, G. *et al.* Tools for probing local circuits: high-density silicon probes combined with optogenetics. *Neuron* **86**, 92–105 (2015).
110. Walter, W. G. *et al.* Contingent Negative Variation : An Electric Sign of Sensori-Motor Association and Expectancy in the Human Brain. *Nature* **203**, 380–384 (1964).
111. Salazar, R. F. *et al.* Content-specific fronto-parietal synchronization during visual working memory. *Science (80-.)*. **338**, 1097–100 (2012).
112. Canolty, R. T. *et al.* High gamma power is phase-locked to theta oscillations in human neocortex. *Science (80-.)*. **313**, 1626–8 (2006).
113. Jensen, O. & Lisman, J. E. Position reconstruction from an ensemble of hippocampal place cells: contribution of theta phase coding. *J Neurophysiol* **83**, 2602–9 (2000).
114. Agarwal, G. *et al.* Spatially distributed local fields in the hippocampus encode rat position. *Science (80-.)*. **344**, 626–30 (2014).
115. Siegel, M., Warden, M. R. & Miller, E. K. Phase-dependent neuronal coding of objects in short-term memory. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 21341–6 (2009).
116. Dean, H. L., Hagan, M. A. & Pesaran, B. Only Coherent Spiking in Posterior Parietal Cortex Coordinates Looking and Reaching. *Neuron* **73**, 829–841 (2012).
117. Pesaran, B., Nelson, M. J. & Andersen, R. A. Free choice activates a decision circuit between frontal and parietal cortex. *Nature* **453**, 406–9 (2008).
118. Hawellek, D. J., Wong, Y. T. & Pesaran, B. Temporal coding of reward-guided choice in the posterior parietal cortex. *Proc Natl Acad Sci USA* (2016). doi:10.1073/pnas.1606479113
119. Vinck, M. *et al.* Gamma-phase shifting in awake monkey visual cortex. *J Neurosci* **30**, 1250–7 (2010).
120. Hastie, T., Tibshirani, R. J. & Friedman, J. *Elements of statistical learning: Data mining, inference and prediction.* (Springer, 2009).
121. Markowitz, D. A., Wong, Y. T., Gray, C. M. & Pesaran, B. Optimizing the decoding of movement goals from local field potentials in macaque cortex. *J Neurosci* **31**, 18412–22 (2011).
122. Bosman, C. A. *et al.* Attentional stimulus selection through selective synchronization between monkey visual areas. *Neuron* **75**, 875–88 (2012).
123. Womelsdorf, T., Fries, P., Mitra, P. P. & Desimone, R. Gamma-band synchronization in visual cortex predicts speed of change detection. *Nature* **439**, 733–6 (2006).
124. Richter, C. G., Thompson, W. H., Bosman, C. A. & Fries, P. A jackknife approach to quantifying single-trial correlation between covariance-based metrics undefined on a single-trial basis. *Neuroimage* **114**, 57–70 (2015).
125. Fröhlich, F. & McCormick, D. A. Endogenous electric fields may guide neocortical network activity. *Neuron* **67**, 129–143 (2010).
126. Anastassiou, C. A. & Koch, C. Ephaptic coupling to endogenous electric field activity: why bother? *Curr Opin Neurobiol* **31**, 95–103 (2015).
127. Cannon, J. *et al.* Neurosystems: brain rhythms and cognitive processing. *Eur J Neurosci* **39**, 705–19 (2014).
128. Besserve, M., Lowe, S. C., Logothetis, N. K., Schölkopf, B. & Panzeri, S. Shifts of Gamma Phase across Primary Visual Cortical Sites Reflect Dynamic Stimulus-Modulated Information Transfer. *PLoS Biol* **13**, e1002257 (2015).
129. Katz, L. N., Yates, J. L., Pillow, J. W. & Huk, A. C. Dissociated functional significance of decision-related activity in the primate dorsal stream. *Nature* **535**, 285–8 (2016).
130. Pesaran, B. & Freedman, D. J. Where Are Perceptual Decisions Made in the Brain? *Trends Neurosci* (2016). doi:10.1016/j.tins.2016.08.008

Investigating large-scale brain dynamics

131. Pettersen, K. H., Hagen, E. & Einevoll, G. T. Estimation of population firing rates and current source densities from laminar electrode recordings. *J Comput Neurosci* **24**, 291–313 (2008).
132. Głąbska, H. T. *et al.* Generalized Laminar Population Analysis (gLPA) for Interpretation of Multielectrode Data from Cortex. *Front Neuroinform* **10**, 1 (2016).
133. Tian, L., Akerboom, J., Schreiter, E. R. & Looger, L. L. Neural activity imaging with genetically encoded calcium indicators. in 79–94 (2012). doi:10.1016/B978-0-444-59426-6.00005-7
134. Chen, F., Tillberg, P. W. & Boyden, E. S. Expansion microscopy. *Science (80-.)*. **347**, 543–548 (2015).
135. Chung, K. *et al.* Structural and molecular interrogation of intact biological systems. *Nature* **497**, 332–337 (2013).
136. Chao, Z. C., Nagasaka, Y. & Fujii, N. Long-term asynchronous decoding of arm motion using electrocorticographic signals in monkeys. *Front Neuroeng* **3**, 3 (2010).
137. Insanally, M. *et al.* A low-cost, multiplexed μ ECoG system for high-density recordings in freely moving rodents. *J Neural Eng* **13**, 026030 (2016).
138. Khodagholy, D. *et al.* NeuroGrid: recording action potentials from the surface of the brain. *Nat. Neurosci.* **18**, 310–5 (2014).
139. Shepherd, G. M. G., Stepanyants, A., Bureau, I., Chklovskii, D. & Svoboda, K. Geometric and functional organization of cortical circuits. *Nat Neurosci* **8**, 782–790 (2005).
140. Lindén, H., Pettersen, K. H. & Einevoll, G. T. Intrinsic dendritic filtering gives low-pass power spectra of local field potentials. *J Comput Neurosci* **29**, 423–44 (2010).
141. Ness, T. V, Remme, M. W. H. & Einevoll, G. T. Active subthreshold dendritic conductances shape the local field potential. *J Physiol* **594**, 3809–3825 (2016).
142. Kajikawa, Y. & Schroeder, C. E. How local is the local field potential? *Neuron* **72**, 847–58 (2011).
143. Nicholson, C. & Llinás, R. Real time current source-density analysis using multi-electrode array in cat cerebellum. *Brain Res* **100**, 418–24 (1975).
144. Pettersen, K. H., Devor, A., Ulbert, I., Dale, A. M. & Einevoll, G. T. Current-source density estimation based on inversion of electrostatic forward solution: effects of finite extent of neuronal activity and conductivity discontinuities. *J Neurosci Meth* **154**, 116–33 (2006).
145. Potworowski, J., Jakuczun, W., Łęski, S. & Wójcik, D. Kernel current source density method. *Neural Comput* **24**, 541–75 (2012).
146. Pettersen, K. H., Lindén, H., Dale, A. M. & Einevoll, G. T. Extracellular spikes and CSD. in *Handbook of Neural Activity Measurement* (eds. Brette, R. & Destexhe, A.) 92–135 (Cambridge University Press, 2012). doi:10.1017/CBO9780511979958.004
147. Rall, W. & Shepherd, G. M. Theoretical reconstruction of field potentials and dendrodendritic synaptic interactions in olfactory bulb. *J Neurophysiol* **31**, 884–915 (1968).
148. Holt, G. R. & Koch, C. Electrical interactions via the extracellular potential near cell bodies. *J Comput Neurosci* **6**, 169–84 (1999).
149. Mazzoni, A. *et al.* Computing the Local Field Potential (LFP) from Integrate-and-Fire Network Models. *PLoS Comp Biol* **11**, e1004584 (2015).
150. Tuckwell, H. C. *Introduction to Theoretical Neurobiology: Volume 1, Linear Cable Theory and Dendritic Structure*. (Cambridge University Press, 1988).
151. Hanes, G. *et al.* Effect of Ionic Diffusion on Extracellular Potentials in Neural Tissue. *PLoS Comp Biol* **12**, e1005193 (2016).
152. Lorente de No, R. Analysis of the distribution of the action currents of nerve in volume conductors. *Stud. Rockefeller Inst. Med. Res. Repr.* **132**, 384–477 (1947).
153. Łęski, S., Lindén, H., Tetzlaff, T., Pettersen, K. H. & Einevoll, G. T. Frequency dependence of signal

Investigating large-scale brain dynamics

- power and spatial reach of the local field potential. *PLoS Comp Biol* **9**, e1003137 (2013).
154. Lindén, H. *et al.* Modeling the spatial reach of the LFP. *Neuron* **72**, 859–72 (2011).
 155. Tenke, C. E., Schroeder, C. E., Arezzo, J. C. & Vaughan, H. G. Interpretation of high-resolution current source density profiles: a simulation of sublaminal contributions to the visual evoked potential. *Exp Brain Res* **94**, 183–192 (1993).
 156. Fernandez-Ruiz, A. *et al.* Cytoarchitectonic and Dynamic Origins of Giant Positive Local Field Potentials in the Dentate Gyrus. *J. Neurosci.* **33**, 15518–15532 (2013).
 157. Haider, B., Schulz, D. P. A., Häusser, M. & Carandini, M. Millisecond Coupling of Local Field Potentials to Synaptic Currents in the Awake Visual Cortex. *Neuron* (2016). doi:10.1016/j.neuron.2016.02.034
 158. Okun, M., Naim, A. & Lampl, I. The subthreshold relation between cortical local field potential and neuronal firing unveiled by intracellular recordings in awake rats. *J. Neurosci.* **30**, 4440–8 (2010).
 159. Głąbska, H. *et al.* Independent Components of Neural Activity Carry Information on Individual Populations. *PLoS One* **9**, e105071 (2014).
 160. Minlebaev, M., Colonnese, M., Tsintsadze, T., Sirota, A. & Khazipov, R. Early γ oscillations synchronize developing thalamus and cortex. *Science (80-.)*. **334**, 226–9 (2011).
 161. Saleem, A. B. *et al.* Subcortical Source and Modulation of the Narrowband Gamma Oscillation in Mouse Visual Cortex. *Neuron* **93**, 315–322 (2017).
 162. Welle, C. G. & Contreras, D. Sensory-driven and spontaneous gamma oscillations engage distinct cortical circuitry. *J. Neurophysiol.* **115**, 1821–1835 (2016).
 163. Bastos, A. M., Briggs, F., Alitto, H. J., Mangun, G. R. & Usrey, W. M. Simultaneous recordings from the primary visual cortex and lateral geniculate nucleus reveal rhythmic interactions and a cortical source for γ -band oscillations. *J Neurosci* **34**, 7639–44 (2014).
 164. Livingstone, M. S. Oscillatory firing and interneuronal correlations in squirrel monkey striate cortex. *J Neurophysiol* **75**, 2467–85 (1996).
 165. Swadlow, H. A., Gusev, A. G. & Bezdudnaya, T. Activation of a cortical column by a thalamocortical impulse. *J Neurosci* **22**, 7766–73 (2002).
 166. Steriade, M., Contreras, D., Amzica, F. & Timofeev, I. Synchronization of fast (30–40 Hz) spontaneous oscillations in intrathalamic and thalamocortical networks. *J. Neurosci.* **16**, 2788–808 (1996).
 167. Swadlow, H. A. & Gusev, A. G. The influence of single VB thalamocortical impulses on barrel columns of rabbit somatosensory cortex. *J Neurophysiol* **83**, 2802–13 (2000).
 168. Schomburg, E. W. *et al.* Theta phase segregation of input-specific gamma patterns in entorhinal-hippocampal networks. *Neuron* **84**, 470–85 (2014).
 169. Buzsáki, G. & Schomburg, E. W. What does gamma coherence tell us about inter-regional neural communication? *Nat Neurosci* **18**, 484–489 (2015).
 170. Castelo-Branco, M., Neuenschwander, S. & Singer, W. Synchronization of visual responses between the cortex, lateral geniculate nucleus, and retina in the anesthetized cat. *J Neurosci* **18**, 6395–6410 (1998).