

1 Straw vs. peat as nest-building material - the impact on farrowing 2 duration and piglet mortality in loose-housed sows

3 Ellen Marie Rosvold*†, Inger-Lise Andersen*

4 * *Norwegian University of Life Sciences, Faculty of Biosciences, Department of Animal and Aquacultural
5 Sciences, PO Box 5003, 1432 Ås, Norway*

6 † *Nord University, Faculty of Biosciences and Aquaculture, PO Box 2501, 7729 Steinkjer, Norway*

7 Corresponding author: Ellen Marie Rosvold. E-mail: ellen.m.rosvold@nord.no

8 9 Abstract

10 Provision of nest-building material pre-partum is required by farm animal legislation in Norway, but as
11 some farmers have problems with straw in the slurry system, alternative and finer-grained substrates
12 could be of interest. The aim of the present study was to examine the effects of providing different nest-
13 building materials before farrowing (peat and long-stemmed straw) on farrowing duration and piglet
14 mortality. Sows of Norsvin Landrace x Swedish Yorkshire were loose-housed in individual farrowing
15 pens with their litter throughout the entire lactation period, and provided with peat (n=18), long-
16 stemmed straw (n=17), or controls (n=18), from two days before expected farrowing until farrowing.
17 Parity ranged from 1 to 9 (mean \pm SE: 2.9 ± 2.0), of which 16 were gilts. The farrowing was video
18 recorded, and the farrowing duration registered from first to last piglet born. Dead piglets were subjected
19 to a post mortem.

20 Sows provided with long-stemmed straw pre-partum had shorter farrowing duration compared to sows
21 provided with peat or controls ($P < 0.001$). The percentage of stillborn piglets was also lowest among
22 sows in the straw group compared to the other two groups ($P < 0.001$). Sows of parity ≥ 4 had longer
23 farrowing duration than gilts and sows of parity 2-3 ($P < 0.001$). Farrowing duration ($P < 0.001$) and
24 the percentage of stillborn piglets increased ($P < 0.001$) with an increasing number of total born piglets.
25 Moreover, increased litter size resulted in a higher pre-weaning mortality ($P = 0.016$).

26 The present study shows that provision of nest-building material before farrowing is important for the
27 birth process of the sow as it reduces farrowing duration and percentage of stillborn piglets compared
28 to controls without relevant substrate. Although peat gave better results than the control treatment, our
29 results suggest that long-stemmed straw is a more appropriate nest-building material.

30 Keywords: Nest building, Nest-building material, Loose-housed sows, Farrowing duration, Stillbirth,
31 Piglet mortality

32 Highlights:

- 33 • Farrowing duration was shorter for loose-housed first parity and older sows provided with nest-
34 building material before farrowing.
- 35 • Loose-housed sows provided with long-stemmed straw had lower percentage of stillborn
36 piglets.
- 37 • Nest-building material had no significant effect on live born mortality.

38 1. Introduction

39 Sows are highly motivated to build a nest before farrowing (e.g. Wischner et al., 2009), and previous
40 studies have found that providing nest-building material, such as straw, has a positive effect on nest-
41 building behaviour (e.g. Thodberg et al., 1999; Westin et al., 2015b; Rosvold et al., 2018). When the
42 sows are allowed to go through a process of nest building, stress is reduced (i.e. hormonal changes; e.g.
43 Yun and Valros (2015)), and maternal behaviour improves, which may indirectly affect piglet survival
44 (Cronin and van Amerongen, 1991; Herskin et al., 1998; Thodberg et al., 1999; Andersen et al., 2005;
45 Yun et al., 2013; 2014). Only a few studies have actually documented a direct relationship between
46 nest-building performance and piglet survival (Ocepek and Andersen, 2017). There are also a few
47 studies showing reduced farrowing duration and reduced incidence of stillbirths when the sow is
48 allowed to express nest-building behaviour with access to relevant material (Thodberg et al., 1999;
49 Westin et al., 2015a; 2015b).

50 Even though straw is a material with positive impact, many farmers hesitate to use it due to the risk of
51 bedding falling through the slatted floor and blocking the drains in the slurry system, and because of
52 more labour needed to maintain pen cleanliness. Peat is finer grained with structural similarities to soil,
53 and suitable for rooting, digging and pawing (Studnitz et al., 2007; Vanheukelom et al., 2011), which
54 also are elements of nest-building behaviour. Therefore, peat is of interest as a nest-building material.

55 The aim of the present project was to investigate the effects of two different nest-building materials on
56 farrowing duration and piglet mortality in individually loose-housed sows. We predicted that provision
57 of peat or long-stemmed straw to sows before farrowing would result in a shorter farrowing duration
58 and a lower piglet mortality compared to sows without access to nest-building material. Due to more
59 positive behavioural effect pre-partum of straw vs. peat (i.e. slightly larger variety of nest-building
60 behaviours, more nest building and a lower frequency of stereotypies; Rosvold et al. (2018)), we also
61 predicted stronger effects of straw than peat on farrowing duration and piglet mortality.

62 2. Materials and methods

63 The present study was conducted in accordance with the Norwegian laws and regulations controlling
64 experiments and procedures on live animals (Nara, 2015).

65 2.1. Experimental design

66 During three farrowing batches, 54 sows, kept loose in individual farrowing pens, were randomly (but
67 balanced for parity) assigned to one of three treatment groups differing in nest-building material: peat,
68 long-stemmed straw or control. There were 18 sows in each treatment group and 18 sows in each batch
69 (6 sows per treatment per batch). The sows were video recorded from two days pre-partum until three
70 days afterwards to document the sows' pre-partum nest-building behaviour, farrowing duration, activity
71 and maternal behaviour. Due to abortion by one sow and failure of video recordings during farrowing
72 of two sows, we collected information about litters and piglet mortality in 53 sows, and farrowing
73 duration of 51 sows.

74 2.2. Animals, housing and feeding

75 The study took place at Mære Agricultural College in Steinkjer, Norway, from May to September 2014.
76 The sows were Norsvin Landrace x Swedish Yorkshire, ranging in parity from 1 to 9 (mean \pm SE: 2.9
77 \pm 2.0), of which 16 were gilts. Mean parity in the treatment groups control, peat and long-stemmed
78 straw were 2.9 \pm 0.5, 3.0 \pm 0.5 and 2.9 \pm 0.5 respectively, and mean parity in the three farrowing batches
79 were 2.7 \pm 0.4, 3.3 \pm 0.6, and 2.9 \pm 0.4 respectively. The sows were inseminated with semen from Duroc
80 boars. The sows were moved from group gestation pens (concrete floor covered with wood-shavings,
81 and sows provided with rooting material and roughage daily) to individual farrowing pens at 3 to 4
82 weeks before farrowing. The farrowing pens had an area of 8.2 m², of which 2.9 m² was slatted flooring
83 (Fig. 1) in one farrowing room with four rows of pens (20 pens in total). The different material
84 treatments were distributed in all four pen rows. According to standard practice in Norway, none of the
85 farrowings were artificially induced, and no laxatives were added to the diet prior to farrowing.

86 The farrowing room was insulated, and mechanically ventilated. The room temperature was regulated
87 to 20°C, and the pen creep area was equipped with floor heating kept at 35°C and heat lamps, which
88 were removed when the piglets were one week of age. The indoor air temperature was measured by two
89 temperature loggers (Tinytag, Gemini Data Loggers, Chichester, UK) placed in different parts of the
90 farrowing room. Due to variation in the outdoor temperature, indoor temperature differed between the
91 batches. From one day before the first farrowing until four days after the last farrowing (12 days), the
92 average temperature was 20.0°C (range 16.8°-24.7°C) for the first batch in May, 23.6°C (19.1°-31.8°C)
93 for the second batch in July, and 20.5°C (17.3°-24.1°C) for the third batch at the end of August.

94 The sows had access to natural light through windows. Consequently, during the summer, it remained
95 light indoors through most of the night. Room lights were on throughout the working day, and only
96 switched on during the night for additional visibility if needed when assisting sows during farrowing.
97 Before farrowing, the sows were fed twice a day by automatic distribution with a lactation concentrate
98 (FK FORMAT Laktasjon, Felleskjøpet, Steinkjer, NO) at approximately 08:30 and 16:00 h, and once
99 during the day with a farrowing concentrate given by hand (FK FORMAT Fødsel, Felleskjøpet,

100 Steinkjer, NO). In accordance with Norwegian legislation, hay (ca 0.3 kg) was distributed to all the
101 sows once every day through the pre-farrowing and lactation periods. After farrowing and through the
102 whole lactation period the sows were fed four times a day by the automatic distribution, and the first 7
103 days after farrowing once daily by hand.

104 After farrowing, wet straw and litter were replaced with dry litter (wood shavings). The sows were
105 taken out of the pen on the first day after farrowing for getting some physical exercise and weight
106 measurement. All the piglets were tooth grinded and they received iron paste orally within the first 24
107 h after birth (Pluss Jernstarter, 1.5 mL; Felleskjøpet). Peat enriched with iron (Pluss Smågristorv,
108 Felleskjøpet) was also provided to all piglets daily from three days after birth (1/2 L, which increased
109 to 1 L per litter within the first two weeks), and concentrate (FK FORMAT Kvikk, Felleskjøpet,
110 Steinkjer, NO) from three days of age. The male piglets were surgically castrated within the first two
111 weeks after birth. Cross-fostering was done between 12-48 h after farrowing. Litter size in this study is
112 thus defined as: number of live born piglets + piglets fostered on – piglets fostered off. Weaning was
113 done at around 5 weeks of age.

114 2.3. Post mortem examination of dead piglets

115 From farrowing until weaning dead piglets were subjected to a post mortem to determine cause of death,
116 and categorized into stillborn (lungs sink when laid in water), crushed (physical signs of crushing were
117 bruising to the body, fractures, haemorrhage or crushed internal organs), starvation (no milk in
118 stomach), and other causes. Piglets that suffered from injuries or starvation, and not able to survive,
119 were euthanized by the staff.

120 2.4. Distribution of nest-building material

121 Nest-building material was provided from two days before expected farrowing. In the morning, the
122 farrowing pens were cleaned and dry wood shavings provided, which had a function as litter for
123 hygienic purposes (0.8 kg, mainly from spruce, same amount to all pens irrespective of treatment, in
124 accordance with Norwegian legislation). Then either 4 kg of peat (90% peat with added formic acid,
125 acetic acid, potassium sorbate and coal; 75% water content, 7.6% crude fiber, and 2.4% ash; Fossli AS,
126 Frosta, NO) or 2 kg of long-stemmed straw (barley) were added to peat and straw treatment pens
127 respectively. Because peat was only about half the volume of straw, the amount was doubled to even
128 out this difference. Sows in the control group did not receive any more material for nest building
129 (Rosvold et al., 2018). In the afternoon the procedure was repeated, with a new provision of litter (0.8
130 kg wood shavings) to each pen if necessary to replace wet and dirty litter, and a refill of 2 kg peat or 1
131 kg long-stemmed straw to the respective treatment groups. The pen cleaning procedure with provision
132 of new litter was done every day until farrowing. Refills of peat were repeated each morning and
133 afternoon until farrowing, as the peat was spread out in the pen because of wallowing and rooting, and

134 disappeared as it was eaten by the sow and went through the slatted floor. Further refill of straw was
135 only necessary if the sow's farrowing was later than expected and dirty straw needed to be replaced.

136 2.5. Video recordings

137 A video camera sensitive to low light (Foscam F19821, 1280x720, Shenzhen, PRC), was suspended
138 above each farrowing pen and connected to a standard PC to record the farrowing process. Farrowing
139 was defined as started when the first piglet was expelled and ended with the birth of the last piglet.

140 2.6. Statistical analysis

141 SAS Version 9.4 (SAS Institute, Inc., Cary, NC) was used to perform statistical analyses. The effects
142 of nest-building material (Control, Peat, Straw), parity (1, 2-3, ≥ 4), batch (1,2,3), litter size (continuous
143 variable), and the interaction between material and parity and the interaction between material and
144 batch, were analysed by a general linear model (PROC GLM) for the following, normally distributed
145 variables: live born, litter size, mortality of live born, percentage of crushed piglets. A generalized
146 model (PROC GENMOD) with Poisson distribution for not normally distributed variables (farrowing
147 duration, stillborn, weaned, starvation and other causes) were also used. Descriptive statistics were
148 obtained using SPSS Version 23 (IBM Corp., Armonk, NY).

149 3. Results

150 3.1. Farrowing duration

151 Mean farrowing duration was 349.9 ± 34.0 min (mean \pm SE), ranging from 90-1235 min (n=51).
152 Provision of straw resulted in significantly shorter farrowing duration than peat, whereas the longest
153 was observed in the control group (Table 1). Farrowing duration increased with parity, as gilts (n=16)
154 spent 262.6 ± 48.7 min on the birth process, sows of parity 2-3 (n=20) spent 362.1 ± 52.4 min, and
155 parity ≥ 4 sows (n=17) 422.9 ± 71.6 min ($\chi^2_{2,35} = 810.75$, $P < 0.001$).

156 There was a significant interaction between material and parity (Table 1, Fig. 2), where sows in the
157 control group of parity ≥ 4 had the longest farrowing duration, and gilts in the straw group had the
158 shortest. Farrowing duration was quite similar between sows in the peat and straw group in parity 1 and
159 ≥ 4 , but a little longer in the latter group, and among sows in parity 2-3, those that received peat had the
160 longest duration.

161 Increased number of total born piglets (live born + stillborn) increased the farrowing duration ($\chi^2_{1,35} =$
162 226.83 , $P < 0.001$; Fig. 5a). Farrowing duration differed between the three batches (mean \pm SE), 359.8
163 ± 57.1 min, 257.6 ± 27.6 min, and 432.4 ± 76.5 min respectively ($\chi^2_{2,35} = 765.07$, $P < 0.001$), and sows
164 in batch 3 had the longest duration. There was also a significant interaction between material and batch
165 on farrowing duration (Table 1). In the control group the farrowing duration (mean \pm SE) in the three
166 farrowing batches was 272.6 ± 79.6 min, 306.7 ± 40.6 min and 761.4 ± 186.5 respectively, in the peat

167 group 451.5 ± 118.5 min, 231.0 ± 51.9 min and 285.7 ± 14.7 respectively, and in the straw group 340.8
168 ± 90.9 min, 230.6 ± 52.3 min and 305.0 ± 64.1 min respectively.

169 3.2. Piglet mortality

170 3.2.1. Overall production and piglet mortality results

171 The 53 sows gave birth to 845 piglets in total; 15.9 ± 0.5 (mean \pm SE) total born per litter (ranging from
172 6 to 24). Of these 794 were live born (15.0 ± 0.5) and 51 stillborn (0.9 ± 0.2). Stillbirths occurred in 28
173 litters (52.8%), and the mean percentage of stillborn piglets (% of total born) was $5.7 \pm 1.0\%$ (ranging
174 from 0-28.6%).

175 The total mortality of live born piglets (% of litter size) was $18.0 \pm 1.9\%$, ranging from 0-55%, and 14.2
176 $\pm 1.7\%$ of the piglets died of crushing, $2.8 \pm 0.8\%$ of starvation and $1.0 \pm 0.3\%$ of other causes. Within
177 the first three days after farrowing, 68.6% of the mortality had occurred. Total number of weaned piglets
178 per sow was 12.0 ± 0.3 (ranging from 6-15).

179 In 20 out of 53 litters (37.7%), the mortality of live born piglets was $>20\%$ (ranging from 21.1% to
180 55.0%). In 14 litters (26.4%) the total mortality of live born piglets was $<8\%$, and in eight litters there
181 was no mortality (Fig. 3).

182 3.2.2. Effects of nest-building materials and parity on piglet mortality

183 Sows provided with straw had the lowest percentage of stillborn piglets, significantly lower than peat
184 and controls, and sows provided with peat had lower percentage of stillborn piglets than controls (Table
185 1). Nest-building material had no significant influence on total mortality of live born piglets or on the
186 different mortality causes. The number of live born piglets, litter size and the number of piglets at
187 weaning were not significantly different between the treatments.

188 Parity had no effect on stillbirths, total piglet mortality, any of the mortality causes or number of weaned
189 piglets. The number of live born piglets increased with increasing parity; Parity 1: 13.1 ± 0.7 , Parity 2-
190 3: 15.7 ± 0.7 , Parity ≥ 4 : 16.0 ± 0.9 ($F_{2,14} = 4.70$, $P = 0.015$), and also litter size; Parity 1: 13.1 ± 0.7 ,
191 Parity 2-3: 15.5 ± 0.6 , Parity ≥ 4 : 15.8 ± 0.9 ($F_{2,14} = 4.72$, $P = 0.015$).

192 There was an interaction between material and parity regarding percentage of stillborn piglets (Table 1,
193 Fig. 4). Sows of parity ≥ 4 provided with long-stemmed straw pre-partum had the lowest percentage of
194 stillborn piglets, whereas first parity sows in the control group had the highest (Fig. 4). There was also
195 a tendency for an interaction between material and parity on mortality of live born piglets (Table 1).

196 3.2.3. Litter size and piglet mortality

197 Increased number of total born piglets resulted in increased percentage of stillborn piglets ($\chi^2_{1,37} = 21.74$,
198 $P < 0.001$; Fig. 5b). The litter size (live born \pm cross-fostered) was 14.9 ± 0.4 piglets (6-24), and
199 increased litter size resulted in increased mortality of live born piglets ($F_{1,15} = 6.37$, $P = 0.016$; Fig. 5c),

200 and a tendency to more crushed piglets ($F_{1,15} = 3.35$, $P = 0.075$). Starvation and other causes of death
201 were not significantly influenced by litter size. Increased litter size resulted in a higher number of
202 weaned piglets ($\chi^2_{1,37} = 8.96$, $P = 0.003$).

203 *3.2.4. Batch and piglet mortality*

204 Total born piglets in the three farrowing batches were 15.9 ± 0.8 , 14.8 ± 0.9 and 17.1 ± 0.9 respectively,
205 not significantly different. Farrowing batch had a significant effect on percentage of stillborn piglets
206 (mean % \pm SE), highest in the first batch; batch 1: $6.6 \pm 1.8\%$, batch 2: $4.7 \pm 1.9\%$, batch 3: $5.7 \pm 1.4\%$
207 ($\chi^2_{2,37} = 17.34$, $P < 0.001$). An interaction effect between material and batch on stillbirths was found
208 (Table 1). In the control group the percentage of stillborn piglets (mean \pm SE) in the three farrowing
209 batches was $9.5 \pm 4.3\%$, $4.7 \pm 2.3\%$ and $10.2 \pm 3.0\%$ respectively, in the peat group $5.6 \pm 2.6\%$, $7.5 \pm$
210 4.6% and $4.8 \pm 1.9\%$ respectively, and in the straw group $4.7 \pm 2.2\%$, $1.3 \pm 1.3\%$ and $2.1 \pm 1.4\%$
211 respectively.

212 There were significant differences in live born mortality between the three batches, $13.7 \pm 2.9\%$, 22.2
213 $\pm 4.0\%$, and $18.4 \pm 2.7\%$ respectively ($F_{2,15} = 3.99$, $P = 0.027$), highest in the second batch, and
214 significant differences regarding crushing in the three batches, $10.5 \pm 2.7\%$, $18.7 \pm 3.7\%$, and $13.8 \pm$
215 2.4% respectively ($F_{2,15} = 3.57$, $P = 0.038$), also highest in the second batch. With respect to starvation,
216 other causes of death and number of weaned piglets there were no significant differences.

217 **4. Discussion**

218 Nest building is a part of the sow's preparation for motherhood. Although nest-building behaviour is
219 largely motivated by internal processes, relevant external stimuli, such as straw, provided at least from
220 around 12 hours pre-partum (Castrén et al., 1993; Andersen et al., 2005), will encourage the sows to
221 spend more time on nest building and express a larger variety of behavioural elements (Westin et al.,
222 2015b; Rosvold et al., 2018). As predicted, the presence of nest-building material resulted in a lower
223 percentage of stillborn piglets compared to the control group, in accordance with previous studies
224 (Westin et al., 2015a), but the effects of material differed between sows of different parity. Percentage
225 of stillborn piglets were lowest for multiparous sows provided with long-stemmed straw, whereas sows
226 in the peat and control groups had a quite similar percentage of stillborn piglets, suggesting that straw
227 is a better material for nest building in experienced sows. However, for first parity sows, the percentage
228 of stillborn piglets did not differ much between the peat and the straw treatment, suggesting that type
229 of material is less crucial for stillbirths in these sows.

230 As predicted, farrowing duration was shorter for sows provided with nesting material than sows in the
231 control group. However, the effects of material was not significant for sows of parity 2 and 3, and the
232 reason for this is unclear. A large number of different studies have documented an association between
233 farrowing duration and the probability of stillbirths (Borges et al., 2005; van Dijk et al., 2005; Canario

234 et al., 2006; Oliviero et al., 2010; Björkman et al., 2017; Thorsen et al., 2017). The risk of hypoxia is
235 higher during a prolonged farrowing, because of damage or occlusion of the umbilical cord, or a
236 placental detachment, which in turn gives a higher risk for stillbirths (van Dijk et al., 2006). Oxytocin
237 release is of importance for effective contractions of the uterus and rapid birth of the piglets (Algers
238 and Uvnäs-Moberg, 2007), and the process of nest building in sows results in elevation of oxytocin
239 levels and a reduction of stress hormones (Yun et al., 2013). It is likely that this is the mechanism
240 explaining why we have these effects on stillbirths and farrowing duration. The control sows had the
241 longest mean farrowing duration, and it is possible that the lack of opportunity to fulfil the behavioural
242 need of nesting induced stress, which in turn led to a prolonged farrowing. These sows also had the
243 highest frequency of stereotypies before farrowing (Rosvold et al., 2018), indicating more frustration
244 and stress.

245 In accordance with some previous studies (Cronin et al., 1993; Björkman et al., 2017), farrowing lasted
246 longer in sows of higher parities in the present study. Aging of the uterus in older sows may reduce
247 muscle tone (Marchant et al., 2000; Borges et al., 2005; Canario et al., 2006) and lead to less efficient
248 expulsions and prolong the farrowing process. Another explanation could be that the farrowing duration
249 also increased with a larger number of piglets born, as seen in earlier research (Rens and Lende, 2004;
250 van Dijk et al., 2005), and sows of higher parities gave birth to larger litters in the present study.

251 In the present study, provision of nest-building material did not affect survival of live born piglets in
252 contrast to a recent study by Swan et al. (2018), and to previous work where a high level of nest-building
253 behaviour pre-partum was associated with lower piglet mortality (Cronin and van Amerongen, 1991),
254 and with no crushing (Andersen et al., 2005). It is possible that the effect on mortality is more dependent
255 on how much time the sows spend on nest building, as loose-housed sows with a high score for nest-
256 building activity had a lower mortality of live born piglets and a higher number of piglets weaned
257 (Ocepek and Andersen, 2017; Ocepek et al., 2017b). Effects of nest-building material are also likely to
258 depend on the amount and the timing of provision. In the present study, we had a limited amount of
259 material. It is possible that free access to nest-building material, with individual needs taken into
260 account, would have given stronger effects, as other studies used ad libitum access from a hayrack the
261 last 24 to 48 hours before expected parturition (e.g. Andersen et al., 2005; Ocepek and Andersen, 2017).
262 Use of hayrack can simplify management of straw as the sow can pull out the amount she needs without
263 spoiling too much on the floor. Parity had no significant effect on mortality of live born piglets in the
264 present study, in accordance with a few studies (Knol et al., 2002; Carney-Hinkle et al., 2013), while
265 others have documented the opposite (e.g. Marchant et al., 2000; Weber et al., 2009; Andersen et al.,
266 2011; Westin et al., 2015a). It is difficult to explain the divergent results on live born mortality with
267 respect to parity, other than the causes are highly multifactorial.

268 A higher number of piglets born increased the percentage of stillborn piglets in the present study, which
269 agrees with previous work (Marchant et al., 2000; Borges et al., 2005; Canario et al., 2006). According
270 to Herpin et al. (2001), larger litters are associated with a longer farrowing duration and a higher risk
271 of hypoxia. Live born piglet mortality and the incidence of crushing (tendency) increased with
272 increasing litter size, in accordance with previous work (Weary et al., 1998; Marchant et al., 2000;
273 Pedersen et al., 2006; Weber et al., 2009; Andersen et al., 2011; Ocepek et al., 2017a). Consistent with
274 earlier work (e.g. Andersen et al., 2011; Kielland et al., 2018), the main cause of death in the present
275 study was crushing. Crushing often occurs immediately after the sow has nosed or oriented towards the
276 piglets, and therefore being aware of their presence. Maternal infanticide by crushing is thus not
277 accidental in many cases, but related to a less protective mothering style (Andersen et al., 2005), and
278 can be considered as a strategy of litter reduction in pigs, especially in larger litters (Andersen et al.,
279 2011; Ocepek et al., 2017a). High litter size also results in more piglets failing to access a teat during a
280 nursing bout, due to a higher sibling competition in larger litters, which is a risk for starvation-related
281 piglet mortality (Andersen et al., 2011; Ocepek et al., 2017a).

282 Farrowing duration was longer in the third and first batch. It is possible that a higher number of total
283 born piglets in these batches was a contributing part of the result. Live born mortality and crushing were
284 higher in the second batch when the temperature was substantially higher than in the other two. More
285 piglets tend to be crushed during the hot season (Weber et al., 2009). Moreover, when surrounding
286 temperatures are above the sow's upper critical temperatures, her appetite is reduced, with a negative
287 consequence for milk production (Quiniou and Noblet, 1999; Cabezón et al., 2017), which is in turn
288 negative for piglet performance (Silva et al., 2018).

289 The mortality of live born piglets was 18% in the present study, in accordance with Kielland et al.
290 (2018) of 18.2%, and slightly lower than the results from Westin et al. (2015a) of 19.5%. The result is
291 higher than the national average in 2014 of 14.2% (Ingris, 2015), and may be explained by the higher
292 number of live born piglets in our study compared to the national average; 15.0 vs. 13.2 piglets (Ingris,
293 2015). Almost 70% of the mortality occurred within the first three days after birth, corresponding with
294 previous studies (e.g. Marchant et al., 2000; Westin et al., 2015a; Kielland et al., 2018).

295 5. Conclusions

296 The results of our study showed that provision of long-stemmed straw or peat to loose-housed sows
297 before farrowing resulted in a shorter farrowing duration compared to controls in sows of most parities.
298 The percentage of stillborn piglets were lower for sows provided with long-stemmed straw compared
299 to both peat and control. The results in the present study thus strengthens the conclusion that long-
300 stemmed straw is best suited as nest-building material.

301 Conflict of interest

302 The authors have no conflicts of interest to declare.

303 Acknowledgements

304 The authors wishes to thank staff at the pig house at Mære Landbruksskole, Heidi Elise Gaundal, Kine
305 Anita Letnes and Geir Næss. The Norwegian Research Council, Norsvin, Animalia, Nortura and Fossli
306 AS (NFR207804/O99) financed this project.

307 References

- 308 Algers, B., Uvnäs-Moberg, K., 2007. Maternal behavior in pigs. *Hormones and Behavior* 52, 78-85.
- 309 Andersen, I.L., Berg, S., Bøe, K.E., 2005. Crushing of piglets by the mother sow (*Sus scrofa*)—purely
310 accidental or a poor mother? *Applied Animal Behaviour Science* 93, 229-243.
- 311 Andersen, I.L., Nævdal, E., Bøe, K.E., 2011. Maternal investment, sibling competition, and offspring
312 survival with increasing litter size and parity in pigs (*Sus scrofa*). *Behav Ecol Sociobiol* 65, 1159-1167.
- 313 Björkman, S., Oliviero, C., Rajala-Schultz, P.J., Soede, N.M., Peltoniemi, O.A.T., 2017. The effect of
314 litter size, parity and farrowing duration on placenta expulsion and retention in sows.
315 *Theriogenology* 92, 36-44.
- 316 Borges, V.F., Bernardi, M.L., Bortolozzo, F.P., Wentz, I., 2005. Risk factors for stillbirth and foetal
317 mummification in four Brazilian swine herds. *Preventive Veterinary Medicine* 70, 165-176.
- 318 Cabezón, F.A., Schinckel, A.P., Marchant-Forde, J.N., Johnson, J.S., Stwalley, R.M., 2017. Effect of
319 floor cooling on late lactation sows under acute heat stress. *Livestock Science* 206, 113-120.
- 320 Canario, L., Cantoni, E., Bihan, E.I., Caritez, J.C., Billon, Y., Bidanel, J.P., Foulley, J.L., 2006. Between-
321 breed variability of stillbirth and its relationship with sow and piglet characteristics. *Journal of*
322 *Animal Science* 84, 3185-3196.
- 323 Carney-Hinkle, E.E., Tran, H., Bundy, J.W., Moreno, R., Miller, P.S., Burkey, T.E., 2013. Effect of dam
324 parity on litter performance, transfer of passive immunity, and progeny microbial ecology¹. *Journal*
325 *of Animal Science* 91, 2885-2893.
- 326 Castrén, H., Algers, B., de Passillé, A.M., Rushen, J., Uvnäs-Moberg, K., 1993. Preparturient variation
327 in progesterone, prolactin, oxytocin and somatostatin in relation to nest building in sows. *Applied*
328 *Animal Behaviour Science* 38, 91-102.
- 329 Cronin, G.M., Schirmer, B.N., McCallum, T.H., Smith, J.A., Butler, K.L., 1993. The effects of providing
330 sawdust to pre-parturient sows in farrowing crates on sow behaviour, the duration of parturition
331 and the occurrence of intra-partum stillborn piglets. *Applied Animal Behaviour Science* 36, 301-315.
- 332 Cronin, G.M., van Amerongen, G., 1991. The effects of modifying the farrowing environment on sow
333 behaviour and survival and growth of piglets. *Applied Animal Behaviour Science* 30, 287-298.
- 334 Herpin, P., Hulin, J.C., Le Dividich, J., Fillaut, M., 2001. Effect of oxygen inhalation at birth on the
335 reduction of early postnatal mortality in pigs. *Journal of animal science* 79, 5-10.
- 336 Herskin, M.S., Jensen, K.H., Thodberg, K., 1998. Influence of environmental stimuli on maternal
337 behaviour related to bonding, reactivity and crushing of piglets in domestic sows. *Applied Animal*
338 *Behaviour Science* 58, 241-254.
- 339 Ingris, 2015. *Årsstatistikk 2014 (Annual Report)*, Animalia Norsvin, Hamar.
- 340 Kielland, C., Wisløff, H., Valheim, M., Fauske, A.K., Reksen, O., Framstad, T., 2018. Preweaning
341 mortality in piglets in loose-housed herds: etiology and prevalence. *animal*, 1-8.
- 342 Knol, E.F., Ducro, B.J., van Arendonk, J.A.M., van der Lende, T., 2002. Direct, maternal and nurse sow
343 genetic effects on farrowing-, pre-weaning- and total piglet survival. *Livestock Production Science*
344 73, 153-164.

345 Marchant, J.N., Rudd, A.R., Mendl, M.T., Broom, D.M., Meredith, M.J., Corning, S., Simmins, P.H.,
346 2000. Timing and causes of piglet mortality in alternative and conventional farrowing systems.
347 Veterinary Record 147, 209-214.

348 Nara, 2015. Regulations on Animal Experiment, [https://lovdata.no/dokument/SF/forskrift/2015-06-](https://lovdata.no/dokument/SF/forskrift/2015-06-18-761)
349 [18-761](https://lovdata.no/dokument/SF/forskrift/2015-06-18-761).

350 Ocepek, M., Andersen, I.L., 2017. What makes a good mother? Maternal behavioural traits
351 important for piglet survival. Applied Animal Behaviour Science 193, 29-36.

352 Ocepek, M., Newberry, R.C., Andersen, I.L., 2017a. Trade-offs between litter size and offspring
353 fitness in domestic pigs subjected to different genetic selection pressures. Applied Animal Behaviour
354 Science 193, 7-14.

355 Ocepek, M., Rosvold, E.M., Andersen-Ranberg, I., Andersen, I.L., 2017b. Can we improve maternal
356 care in sows? Maternal behavioral traits important for piglet survival in loose-housed sow herds1.
357 Journal of Animal Science 95, 4708-4717.

358 Oliviero, C., Heinonen, M., Valros, A., Peltoniemi, O., 2010. Environmental and sow-related factors
359 affecting the duration of farrowing. Animal Reproduction Science 119, 85-91.

360 Pedersen, L.J., Jørgensen, E., Heiskanen, T., Damm, B.I., 2006. Early piglet mortality in loose-housed
361 sows related to sow and piglet behaviour and to the progress of parturition. Applied Animal
362 Behaviour Science 96, 215-232.

363 Quiniou, N., Noblet, J., 1999. Influence of high ambient temperatures on performance of
364 multiparous lactating sows. Journal of animal science 77, 2124-2134.

365 Rens, B.T.T.M.v., Lende, T.v.d., 2004. Parturition in gilts: duration of farrowing, birth intervals and
366 placenta expulsion in relation to maternal, piglet and placental traits. Theriogenology 62, 331-352.

367 Rosvold, E.M., Newberry, R.C., Framstad, T., Andersen, I.-L., 2018. Nest-building behaviour and
368 activity budgets of sows provided with different materials. Applied Animal Behaviour Science 200,
369 36-44.

370 Silva, B.A.N., Tolentino, R.L.S., Eskinazi, S., Jacob, D.V., Raidan, F.S.S., Albuquerque, T.V., Oliveira,
371 N.C., Araujo, G.G.A., Silva, K.F., Alcici, P.F., 2018. Evaluation of feed flavor supplementation on the
372 performance of lactating high-prolific sows in a tropical humid climate. Animal Feed Science and
373 Technology 236, 141-148.

374 Studnitz, M., Jensen, M.B., Pedersen, L.J., 2007. Why do pigs root and in what will they root?: A
375 review on the exploratory behaviour of pigs in relation to environmental enrichment. Applied Animal
376 Behaviour Science 107, 183-197.

377 Swan, K.-M., Peltoniemi, O.A.T., Munsterhjelm, C., Valros, A., 2018. Comparison of nest-building
378 materials in farrowing crates. Applied Animal Behaviour Science.

379 Thodberg, K., Jensen, K.H., Herskin, M.S., Jørgensen, E., 1999. Influence of environmental stimuli on
380 nest building and farrowing behaviour in domestic sows. Applied Animal Behaviour Science 63, 131-
381 144.

382 Thorsen, C.K., Schild, S.-L.A., Rangstrup-Christensen, L., Bilde, T., Pedersen, L.J., 2017. The effect of
383 farrowing duration on maternal behavior of hyperprolific sows in organic outdoor production.
384 Livestock Science 204, 92-97.

385 van Dijk, A.J., van der Lende, T., Taverne, M.A.M., 2006. Acid–base balance of umbilical artery blood
386 of liveborn piglets at birth and its relation with factors affecting delivery of individual piglets.
387 Theriogenology 66, 1824-1833.

388 van Dijk, A.J., van Rens, B.T.T.M., van der Lende, T., Taverne, M.A.M., 2005. Factors affecting
389 duration of the expulsive stage of parturition and piglet birth intervals in sows with uncomplicated,
390 spontaneous farrowings. Theriogenology 64, 1573-1590.

391 Vanheukelom, V., Driessen, B., Maenhout, D., Geers, R., 2011. Peat as environmental enrichment for
392 piglets: The effect on behaviour, skin lesions and production results. Applied Animal Behaviour
393 Science 134, 42-47.

394 Weary, D.M., Phillips, P.A., Pajor, E.A., Fraser, D., Thompson, B.K., 1998. Crushing of piglets by sows:
395 effects of litter features, pen features and sow behaviour. *Applied Animal Behaviour Science* 61,
396 103-111.

397 Weber, R., Keil, N.M., Fehr, M., Horat, R., 2009. Factors affecting piglet mortality in loose farrowing
398 systems on commercial farms. *Livestock Science* 124, 216-222.

399 Westin, R., Holmgren, N., Hultgren, J., Ortman, K., Linder, A., Algers, B., 2015a. Post-mortem findings
400 and piglet mortality in relation to strategic use of straw at farrowing. *Preventive Veterinary Medicine*
401 119, 141-152.

402 Westin, R., Hultgren, J., Algers, B., 2015b. Strategic use of straw increases nest building in loose
403 housed farrowing sows. *Applied Animal Behaviour Science* 166, 63-70.

404 Wischner, D., Kemper, N., Krieter, J., 2009. Nest-building behaviour in sows and consequences for
405 pig husbandry. *Livestock Science* 124, 1-8.

406 Yun, J., Swan, K.-M., Farmer, C., Oliviero, C., Peltoniemi, O., Valros, A., 2014. Prepartum nest-building
407 has an impact on postpartum nursing performance and maternal behaviour in early lactating sows.
408 *Applied Animal Behaviour Science* 160, 31-37.

409 Yun, J., Swan, K.-M., Vienola, K., Farmer, C., Oliviero, C., Peltoniemi, O., Valros, A., 2013. Nest-
410 building in sows: Effects of farrowing housing on hormonal modulation of maternal characteristics.
411 *Applied Animal Behaviour Science* 148, 77-84.

412 Yun, J., Valros, A., 2015. Benefits of Prepartum Nest-building Behaviour on Parturition and Lactation
413 in Sows - A Review.

414

415 **Figure captions**

416 **Table 1.** Litter traits and mortality causes (mean \pm SE): effects of nest-building material, the interaction
417 between nest-building material and parity, and the interaction between nest-building material and batch.

418 **Fig. 1.** The design of the farrowing pen.

419 **Fig. 2.** The interaction between material and parity regarding farrowing duration, min (mean \pm SE).

420 **Fig. 3.** Frequency distribution of the mortality of live born per sow (%), n=53.

421 **Fig. 4.** The interaction between material and parity regarding stillborn piglets (mean % \pm SE).

422 **Fig. 5.** (a) Farrowing duration (min) in relation to total born piglets (live born + stillborn), n=51. (b)
423 Stillborn piglets (mean %) with respect to total born piglets, n=53 litters. (c) Mortality of live born
424 piglets (%) with respect to litter size (live born \pm cross-fostered), n=53 litters.




Fig. 1. The design of the farrowing pen.




Fig. 2. The interaction between material and parity regarding farrowing duration, min (mean \pm SE).




Fig. 3. Frequency distribution of the mortality of live born per sow (%), n=53.




Fig. 4. The interaction between material and parity regarding stillborn piglets (mean % \pm SE).




Fig. 5. (a) Farrowing duration (min) in relation to total born piglets (live born + stillborn), n=51. (b) Stillborn piglets (mean %) with respect to total born piglets, n=53 litters. (c) Mortality of live born piglets (%) with respect to litter size (live born ± cross-fostered), n=53 litters.

Table 1. Litter traits and mortality causes (mean \pm SE): effects of nest-building material, the interaction between nest-building material and parity, and the interaction between nest-building material and batch.

	Material			$\chi^2_{2,37}$	P-value	Material x Parity		Material x Batch	
	Control (n=18)	Peat (n=18)	Straw (n=17)			$\chi^2_{4,37}$	P-value	$\chi^2_{4,37}$	P-value
Farrowing duration, min	438.1 \pm 82.6 ^a (n=16)	322.7 \pm 46.7 ^b (n=18)	295.8 \pm 41.1 ^c (n=17)	262.72 ¹	<0.001	307.31 ²	<0.001	991.28 ²	<0.001
Stillborn, % ³	8.1 \pm 1.9 ^a	6.0 \pm 1.8 ^b	2.8 \pm 1.0 ^c	44.79	<0.001	39.22	<0.001	20.09	<0.001
Live born	14.4 \pm 0.9	15.8 \pm 1.0	14.7 \pm 0.7	F _{2,14} = 1.06	0.3573	F _{4,14} = 1.09	0.3764	F _{4,14} = 1.94	0.1229
Litter size ⁴	14.4 \pm 0.8	15.4 \pm 0.9	14.8 \pm 0.6	F _{2,14} = 0.72	0.493	F _{4,14} = 0.69	0.601	F _{4,14} = 1.52	0.215
Weaned	11.7 \pm 0.6	11.9 \pm 0.6	12.3 \pm 0.5	0.44	0.803	2.27	0.687	1.26	0.8688
Live born mortality, % ⁵	16.8 \pm 3.8	21.1 \pm 3.1	16.1 \pm 2.8	F _{2,15} = 0.56	0.574	F _{4,15} = 2.27	0.080	F _{4,15} = 1.05	0.395
Crushed, %	12.4 \pm 3.2	16.3 \pm 2.8	14.0 \pm 2.7	F _{2,15} = 0.43	0.654	F _{4,15} = 1.29	0.293	F _{4,15} = 0.98	0.432
Starvation, %	3.3 \pm 1.6	3.7 \pm 1.6	1.4 \pm 0.7	- ⁶	- ⁶	- ⁶	- ⁶	- ⁶	- ⁶
Other causes, %	1.1 \pm 0.6	1.1 \pm 0.6	0.7 \pm 0.5	- ⁶	- ⁶	- ⁶	- ⁶	- ⁶	- ⁶

¹ $\chi^2_{2,35}$

² $\chi^2_{4,35}$

³ % of total born (live born + stillborn).

⁴ Litter size: live born \pm cross-fostered piglets.

⁵ % of litter size.

⁶ Number of observations too low to analyse.

^{a-c} Means with superscripts are significantly different (P < 0.01).