

1 **Farm scale modelling of greenhouse gas emissions from semi-intensive suckler cow beef**
2 **production**

3 Stine Samsonstuen^{a*}, Bente A. Åby^a, Paul Crosson^b, Karen A. Beauchemin^c, Helge Bonesmo^d,
4 Laila Aass^a

5 ^a Department of Animal Sciences, Norwegian University of Life Sciences, P.O. Box 5003 NO-
6 1432 Ås, Norway

7 ^b Animal Grassland Research and Innovation Centre, Teagasc, Grange, Dunsany, Co. Meath,
8 Ireland

9 ^c Lethbridge Research and Development Centre (LeRDC), Agriculture and Agri-Food Canada,
10 Lethbridge, T1J4B1, Canada

11 ^d Felleskjøpet Fôrutvikling, NO-7018 Trondheim, Norway

12 *Corresponding author at: Department of Animal Sciences, Norwegian University of Life
13 Sciences, P.O. Box 5003 NO-1432 Ås, Norway. E-mail address: stine.samsonstuen@nmbu.no
14 (S.Samsonstuen).

15 **Abstract**

16 A whole-farm model, HolosNorBeef was developed to estimate net greenhouse gas (GHG)
17 emissions from suckler beef production systems in Norway. The model considers direct
18 emissions of methane (CH₄), nitrous oxide (N₂O) and carbon dioxide (CO₂) from on-farm
19 livestock production including soil carbon (C) changes, and indirect N₂O and CO₂ emissions
20 associated with leaching, volatilization and inputs used on the farm. The emission intensities
21 from average beef cattle farms in Norway was estimated by considering typical herds of British
22 and Continental breeds located in two different regions, flatlands and mountains, with different
23 resources and quality of feed available. The flatlands was located at a low altitude in an area
24 suitable for grain production and mountains was located at higher altitude in a mountainous area

25 not suitable for grain production. The estimated emission intensities were 29.5 and 32.0 kg CO₂
26 equivalents (eq) kg⁻¹ carcass for the British breeds and 27.5 and 29.6 kg CO₂ eq kg⁻¹ carcass for
27 the Continental breeds, for flatlands and mountains, respectively. Enteric CH₄ was the largest
28 source accounting for 44-48% of total GHG emissions. Nitrous oxide from manure and soil was
29 the second largest source accounting for, on average, 21% of the total emissions. Carbon
30 sequestration reduced the emission intensities by 3% on average. When excluding soil C the
31 difference between locations decreased in terms of GHG emission intensity, indicating that
32 inclusion of soil C change is important when calculating emission intensities, especially when
33 production of feed and use of pasture are included.

34 **Keywords**

35 Beef cattle; greenhouse gas emissions; farm scale model; soil carbon; suckler cow production

36 **1. Introduction**

37 The global population is expected to reach 9.73 billion by 2050 and it is estimated that global
38 food production needs to increase by 50% compared with 2012 levels (FAO, 2017). Human
39 population growth and climate change are exerting pressure on agricultural production systems
40 to secure food production while minimizing greenhouse gas (GHG) emissions. In 2015,
41 agriculture accounted for 10% of the total GHG emissions in Europe (European Environment
42 Agency, 2017). It is a political goal to reduce total GHG emissions 40% by 2030 compared with
43 1990 levels (European Commission, 2014) and the agricultural sector is expected to contribute.

44 In compliance with policy commitments to reducing total GHG emissions, livestock
45 supply chains have focused on decreasing GHG emission intensity, which is a measure of the
46 quantity of GHG emissions generated in the production of a product. Focusing on emission

47 intensity allows the industry to grow, but with less GHG emissions relative to the amount of
48 product produced. In the case of beef, it is necessary to reduce emission intensities considerably,
49 as global beef production is expected to increase by 72% when compared to 2000 levels (FAO,
50 2006). The emission intensity of beef production has been investigated in a number of studies
51 (Beauchemin et al., 2010; Beauchemin et al., 2011; Foley et al., 2011; Mogensen et al., 2015;
52 Alemu et al., 2017) and varies widely, ranging from 17-37 CO₂ eq (kg⁻¹ carcass) and 16.3-38.8
53 CO₂ eq (kg⁻¹ live weight sold). The substantial variation in GHG emissions intensities for beef
54 production systems are due to differences in farming systems (Nguyen et al., 2010), location
55 (White et al., 2010) and farm management (Alemu et al., 2017). In terms of farm management, it
56 has been shown that farm technical efficiency improvements have an important role to play in
57 reducing GHG emissions intensity (Beauchemin et al., 2011; Zhang et al., 2013).

58 Whole farm systems models are useful for assessing the impact of improvements in
59 technical efficiency and direct mitigation options on farm-level GHG emissions and emission
60 intensity. In a review of farm-level modelling approaches by Schils et al. (2007) it was
61 concluded that a whole-farm approach is a powerful tool for development of cost effective
62 mitigation options, as interactions between farm components are revealed.

63 Previous studies have found substantial differences in emission intensities among
64 continents (Gerber et al., 2013) and among farms within a country (Bonesmo et al., 2013),
65 depending upon natural resources and farm management. Norway is a country with varying
66 production conditions, with large areas suitable as pastures and only a small area (1%) suitable
67 for grain production (Åby et al., 2014), limited by climate and topography. Most farm-level
68 modelling studies assume that soil carbon (C) is at equilibrium. However, Soussana et al. (2007)

69 concluded that European grasslands are likely to act as atmospheric C sinks. The net impact of
70 including soil C in farm level modeling studies of beef production is not clear.

71 Thus, the aim of this study was to 1) develop a whole farm GHG model, HolosNorBeef,
72 which includes changes in soil C and is adapted to the various production systems and feed
73 resources in Norway, and 2) to use the model to evaluate the GHG emissions from typical
74 suckler beef cow herds in two geographically different regions of Norway with different
75 resources and quality of feed available.

76 **2. Materials and methods**

77 2.1 HolosNorBeef

78 The HolosNorBeef model was developed to estimate net GHG emissions from suckler beef
79 production systems in Norway. It is an empirical model based on the HolosNor model (Bonesmo
80 et al., 2013), BEEFGEM (Foley et al., 2011) and the methodology of the Intergovernmental
81 Panel on Climate Change (IPCC, 2006) modified for suckler beef production systems under
82 Norwegian conditions. The suckler cow beef production system in Norway is semi-intensive
83 with extensive (low concentrate; approx. 0-10%) feeding of suckler cows, calves and heifer
84 progeny and intensive (high concentrate; approx. 50%) finishing of male progeny as bulls for
85 meat production (Åby et al., 2012). Suckler cows are kept indoors on during winter (approx. 8
86 months) during which time they are fed grass silage, hay or straw and minimal amounts of
87 concentrates. During summer (approx. June to mid-September) they are kept on pasture with
88 their calves. Mating season is during pasture and the calving season is from March to mid-June.
89 Calves are weaned at 6 months of age, and the bull progeny are then fed a high concentrate diet
90 (approx. 50%) until they are slaughtered at a relatively early age (average 16.7 months;

91 Animalia, 2017a). Heifers are retained as replacements, sold or slaughtered. The cow-calf
92 enterprise and finishing of bulls take place at the same farm. The most numerous breeds in
93 Norway are: Charolais, Hereford, Limousin, Aberdeen Angus and Simmental (Animalia, 2017b).
94 Data for the present study were obtained from The Norwegian Beef Cattle Herd Recording
95 System that maintains individual data for animals from birth to slaughter, including weights,
96 reproductive traits and carcass data. HolosNorBeef also includes the data for feed resources,
97 diets and manure management, soil characteristics and weather.

98 HolosNorBeef was developed in Microsoft Excel (Microsoft Corporation, 2016) and is a
99 two-step model where the first sub-model incorporates a detailed description of the farm to be
100 used in the second sub-model (Section 2.1.1) that estimates on-farm GHG emissions (Section
101 2.1.2.) using a cradle to farm gate approach. The GHG sub-model considers direct emissions of
102 methane (CH₄), nitrous oxide (N₂O) and carbon dioxide (CO₂) from on-farm livestock
103 production including soil C changes, and indirect N₂O and CO₂ emissions associated with run-
104 off, nitrate leaching, ammonia volatilization and from inputs used on the farm (Figure 1). Direct
105 emissions from animal production are calculated on a monthly basis, accounting for diet and
106 weather differences. All GHG emissions are expressed as CO₂-equivalents (eq) to account for the
107 global warming potential of the respective gases for a time horizon of 100 years:

108 $CH_4(\text{kg}) \times 28 + N_2O(\text{kg}) \times 265 + CO_2(\text{kg})$ (Myhre et al., 2013). Emissions intensities are
109 expressed as GHG emissions (kg CO₂ eq) per kg beef carcass produced.

110 *2.1.1 Input sub-model*

111 The input sub-model gives a detailed description of the number of animals in each class of cattle,
112 the animal live weights, energy requirements and feed intake on a monthly basis. The monthly

113 live weights for each class of cattle are based on birth weights, weaning weights, yearling
114 weights, slaughter weights and adult weights. The weight at the start of each month are
115 calculated based on the starting live weight and live weight change for the previous month. The
116 number of animals in each class of cattle at the start of each month is based on the number at the
117 start of the previous month adjusted for the number of calvings, stillbirths, twin frequency,
118 mortality rate and any sales and purchases in the previous month. The replacement rate is set to
119 keep the farm size constant and kg beef carcass produced is calculated based on the number of
120 animals sold to abattoirs, slaughter weights and dressing percentages.

121 Daily energy requirements of each class of cattle are estimated according to Refsgaard
122 Andersen (1990) and are based on the animals' requirements for maintenance, growth,
123 pregnancy and lactation. Dry matter intake (DMI) considers the energy requirements of the
124 animal and the animals' intake capacity and is calculated for each animal group. Intake capacity
125 is dependent on the fill value of the forage as well as the substitution rate of the concentrates
126 (Refsgaard Andersen, 1990). Gross energy (GE) intake is estimated based on dry matter intake
127 and the GE content of the diet. The nutrient content of the diet is determined from the chemical
128 composition of commercial concentrates produced by the two largest feed mills in Norway
129 (Felleskjøpet SA, Oslo Norway; Norgesfor AS, Oslo Norway) and forages (laboratory analysis
130 information provided by Eurofins, Moss Norway).

131 *2.1.2 GHG emissions sub-model*

132 *Methane emissions*

133 HolosNorBeef estimates enteric CH₄ emissions for each class of cattle using an IPCC (2006) Tier
134 2 approach. Enteric CH₄ emissions are calculated from GE intake using an adjusted CH₄
135 conversion factor ($Y_m = 0.065$; IPCC 2006). The Y_m is adjusted for the digestibility of the diet

136 according to Bonesmo et al. (2013), as suggested by Beauchemin et al. (2010; Table 1). Manure
137 CH₄ emissions are based on the production of volatile solids (VS) according to IPCC (2006),
138 taking the GE content and digestibility of the diet into account. The VS production is multiplied
139 by a maximum CH₄ producing capacity of the manure ($B_o=0.18 \text{ m}^3 \text{ CH}_4 \text{ kg}^{-1}$) and a CH₄
140 conversion factor specific for the management practice used (Table 1).

141 *Nitrous oxide emissions*

142 The direct N₂O emissions from manure are calculated by multiplying the manure N content with
143 an emission factor for the manure handling system; deep bedding or deposited on pasture (Table
144 1; IPCC, 2006). Manure N content is estimated based on DMI, crude protein (CP; $\text{CP} = 6.25 \times$
145 N) content of the diet and N retention by the animals based on IPCC (2006).

146 Direct N₂O emissions from soils are estimated based on N inputs, using the IPCC (2006)
147 emission factor of $0.01 \text{ kg N}_2\text{O-N kg}^{-1} \text{ N}$ applied. Total N inputs include application of N
148 fertilizer and manure, grass and crop residual N and mineralized N (Table 1). Straw from grain
149 crop is left on the fields and is included in residue N. Residue N is calculated as the sum of
150 above- and below ground residue, using the crop yields of Janzen et al. (2003). Mineralization of
151 N inputs is calculated using the derived C:N ratio of organic soil matter of 0.1 (Little et al.,
152 2008). To account for location specific effects of soil moisture and temperature, the relative
153 effects of percentage water filled pore space (WFPS) of top soil and soil temperature at 30 cm
154 depth ($t_{s30^\circ\text{C}}$) are based on Sozanska et al. (2002) and included as described by Bonesmo et al.
155 (2012; Table 1). Seasonal variations were taken into account by including four seasons; spring
156 (April-May), summer (June-August), fall (September-November) and winter (December-March).
157 The “timing effect” of the application of N fertilizer and manure were calculated using a crop

158 specific factor (Sozanska et al. 2002) and used to calculate the N₂O-N for each season based on
159 WFPS and ts30°C.

160 The indirect N₂O emissions emitted on farm from run-off, leaching and volatilization
161 (Table 1) are estimated from assumed losses of N from manure, residues and fertilizer according
162 to IPCC (2006). The emissions were estimated based on the assumed fraction of N lost adjusted
163 for emission factors (0.0075 and 0.01 kg N₂O-N kg⁻¹) for leaching and volatilized ammonia-N,
164 respectively (IPCC, 2006).

165 *Soil C change*

166 Estimates of soil C change are based on the Introductory Carbon Balance Model (ICBM) by
167 Andrén et al. (2004). The model considers two soil C pools; young (Y) and old (O), accounting
168 for 7% and 93% of the initial C content of the top soil, respectively. The change in Y and O soil
169 C are estimated from total C inputs (i), a humification coefficient (h; Table 1), two decay
170 constants (k_Y and k_O; Table 1) and the relative effect of soil moisture (r_w) and temperature (r_T).
171 Total soil C inputs are calculated from crop residues and manure as described by Andrén et al.
172 (2004). Similar to HolosNor (Bonesmo et al., 2013), regional differences are accounted for by
173 including annual soil and climate data, which are based on the specific crop and soil type
174 together with weather data from specific sites. The yearly C fluxes of Y and O soil C are given
175 by the differential equations of Andrén and Kätterer (1997):

$$176 \quad \frac{dY}{dt} = i - k_1 r Y$$

$$177 \quad \frac{dO}{dt} = h k_1 r Y - k_2 r O$$

178 *Carbon dioxide emissions*

179 HolosNorBeef estimates CO₂ emissions from energy use. Direct emissions from use of diesel
180 fuel and off-farm emissions from production and manufacturing of farm inputs (i.e. fertilizers
181 and pesticides) are estimated using emission factors from Norway or Northern-Europe (Table 1).
182 Indirect emissions related to purchased concentrates are estimated according to Bonesmo et al.
183 (2013). The amount of purchased concentrates is estimated based on the concentrate deficit,
184 determined as the concentrate required to meet the energy and CP requirements minus grain and
185 oilseeds grown on the farm. The deficit is assumed to be supplied by barley and oats grown in
186 Norway and soybean meal imported from South America (Table 1). On-farm emissions from
187 production of field crops produced on the farm but not used in the beef enterprise (e.g. either
188 sold or consumed by other classes of farm animals) are not included in the total farm emissions
189 related to beef production.

190 2.2 Norwegian suckler beef production system

191 Four farms representative of beef production systems in Norway were modelled. The farms
192 represent 'typical' Norwegian farms in term of scale, production results, feeding regimes and
193 location within the country. The locations chosen for the study are areas with a large proportion
194 of Norwegian suckler cow production and are referred to as flatlands and lowlands. The
195 administrative center of flatlands (latitude/longitude 60.9/10.7) has an altitude of 246 m above
196 sea level (m.a.s.l), whereas mountains (latitude/longitude 62.5/9.7) is located at 545 m.a.s.l.. The
197 locations have different resource bases and average temperatures (Table 2), and on a scale from 1
198 (good) to 8 (harsh) as compiled by Norwegian Meterological Insitute and Det norske
199 hageselskap (2006), flatlands and mountains are within climatic zone 4 and 7, respectively. The

200 locations differ in farm size and areas available for forage and crop production, which influence
201 the use of different input factors.

202 The input data were average beef cattle production data (Åby et al., 2012; Animalia,
203 2017a; Animalia, 2017b), farm operational data from the Norwegian Institute of Bioeconomy
204 Research (NIBIO, 2015) and soil and weather data (Skjelvåg et al., 2012) for the specific
205 locations. The farm operational data are annual status reports based on tax results from a
206 representative random sample of 81 Norwegian farms distributed across the country, whereas 21
207 and 11 were located in the flatland and mountains, respectively (NIBIO, 2015). In each location
208 an average herd of British (Angus and Herford) and Continental (Limousin, Simmental and
209 Charolais) breeds were considered. The breed specific weights at different ages, proportion of
210 stillborn calves, twin frequency and proportion dead before 180 days (Table 3) were obtained
211 from Åby et al. (2012), Animalia (2017a) and Animalia (2017b). The herd size and number of
212 cattle in each class were based on average number of cows, average number of calvings and
213 average number of heifers and calves (Table 4) obtained from NIBIO (2015). Estimates of
214 proportion of concentrates and time spent on pasture for each cattle class were available from
215 Åby et al. (2012). The manure was assumed to be deposited on pasture during the grazing period
216 and during housing the manure handling system was deep bedding. The areas (ha) and yields (kg
217 ha⁻¹) of grass, barley, oats, winter wheat and summer wheat were obtained from NIBIO (2015;
218 Table 4). The reduced tillage ratios for oats, barley, spring- and winter wheat were zero. The DM
219 contents and nutritive values of the grass silages were estimated using data from Eurofins for the
220 specific locations (Table 4). Use of energy, fuel and pesticides were available through the costs
221 (NIBIO, 2015; Table 4). Cost of pesticides was distributed to the various crops according to
222 Bonesmo et al. (2013) using relative weighting factors: barley, 1.00; oats, 0.51; spring wheat,

223 1.05; winter wheat, 1.71; and grass production, 0.15. The use of fertilizers was based on the
224 Norwegian recommendations for N, P and K application levels for the specific crops (Table 4).
225 Seasonal soil and weather data were available through Skjelvåg et al. (2012; Table 5).

226 2.3 Sensitivity analysis

227 A sensitivity analysis was performed to evaluate possible errors in the most important emission
228 factors (EF): CH₄ conversion factor (Y_m), manure N₂O (IPCC, 2006), soil N₂O (IPCC, 2006),
229 manufacturing of N-fertilizer (DNV, 2010), and a combined indirect and direct EF for fuel (The
230 Norwegian Environment Agency, 2017; Öko-Institut, 2010). In addition, the sensitivity of the
231 yearly effect of temperature and soil moisture ($r_w \times r_T$), and initial soil organic carbon content
232 was investigated. A farm with British breeds located in the flatlands were chosen as a baseline
233 for the sensitivity analysis. Emission factors were changed one percent, and emission intensities
234 were re-calculated and related to the baseline as a percentage change in emission intensities. The
235 sensitivity of farm and herd size was tested based on variation in the farm operational data from
236 NIBIO (2015) by evaluating a small and a large farm of British breeds located in the flatlands
237 (Table 6).

238 3. Results

239 The total emissions ranged from 227 to 284 t CO₂ eq. In both locations British breeds had less
240 total net emissions than Continental breeds (Table 7). Enteric CH₄, manure CH₄ and manure N₂O
241 emissions were greater for the Continental breeds in both locations. Soil N₂O emissions were
242 greater for flatlands. Flatlands had greater soil C sequestration and greater energy CO₂
243 emissions.

244 Enteric CH₄ contributed most to the GHG emissions, accounting for 44-48% of the
245 emissions (Table 7). Nitrous oxide from manure and soil were the second largest source, each
246 accounting for on average 10% of the total emission. Direct CH₄ emissions from manure
247 accounted for 10-12% of total emissions. Soil C balance was negative for Continental breeds in
248 both locations and British breeds in flatlands, indicating C sequestration. However, British
249 breeds had positive soil C in mountains, indicating a loss of soil C. The on-farm direct emissions
250 from burning of fossil fuels accounted for 5-8% of the total emissions.

251 The emission intensities were greater for the British breeds (29.5 to 32.0 kg CO₂ eq kg⁻¹
252 carcass) compared with the Continental breeds (27.5 to 29.6 kg CO₂ eq kg⁻¹ carcass) in both
253 locations (Table 8).

254 Enteric CH₄ conversion factor had the highest sensitivity elasticity, having a 0.45%
255 change in emission intensities caused by one percentage change in Y_m (Table 9). The estimated
256 GHG were moderate sensitive to changes in manure N₂O EF, soil N₂O EF, N-fertilizer EF, and
257 fuel EF ranging from 0.09 to 0.12%. The initial soil organic carbon and the yearly effect of soil
258 temperature and soil moisture ($r_w \times r_T$) had a moderate linear and moderate non-linear response,
259 respectively (Table 9). The total emissions increased with increasing farm and herd size. In terms
260 of emission intensities, the changed farm and herd size increased the emission intensities for the
261 small farm and reduced the emission the emission intensities for the large farm (Table 10).

262 **4. Discussion**

263 The HolosNorBeef model is derived from IPCC methodology (2006) with modifications to
264 accommodate Norwegian conditions, similar to the original HOLOS model developed for
265 Canada (Little et al., 2008). Most whole-farm system models are based on IPCC methodology

266 (Crosson et al., 2011), but adapting the methodology for local, regional or national conditions
267 improves the sensitivity of the model to differences in production and environmental
268 circumstances. The estimated emission intensities in the present study are comparable with the
269 range of intensities for beef presented by Crosson et al. (2011). The range of emission intensities
270 across studies for different countries and production systems reflects the differences in
271 assumptions, algorithms and approaches in addition to the differences in farm management,
272 breed differences and natural resources. Direct comparisons across studies should therefore be
273 done with caution.

274 The assessment in the present study used a cradle to farm gate approach, simulating both
275 internal and external flows of the input factors to calculate the GHG emissions of beef
276 production (Figure 1). A whole-farm approach ensures that interactions are taken into account,
277 and that the effects of changes in one factor are transferred throughout the system (Schils et al.,
278 2007).

279 HolosNorBeef estimated emission intensities for average herds of British and Continental
280 breeds in Norway of 27.5-32.0 CO₂ eq (kg carcass)⁻¹. This range of intensities is similar to the
281 emission intensities reported for farming systems in Ireland: 23.1 CO₂ eq (kg carcass)⁻¹ (Foley et
282 al., 2011), Denmark: 23.1-29.7 CO₂ eq (kg carcass)⁻¹ and Sweden: 25.4 CO₂ eq (kg carcass)⁻¹
283 (Mogensen et al., 2015). In those studies, emission intensities from enteric CH₄ varied depending
284 upon the on feeding intensity (Ireland, 49.1% of total GHG emissions; Denmark/Sweden, 47.6-
285 55.65% of total GHG emissions). In the present study, enteric CH₄ varied from 43.9-48.2% of
286 total GHG emissions for the two breeds (Table 6). Mitigation strategies are often aimed at
287 reducing enteric CH₄ emissions. The CH₄ conversion factor (i.e. Y_m) had the highest sensitivity
288 elasticity, thus a reliable Y_m is crucial as a significant change in Y_m due to feeding intensity

289 would influence the emission intensities considerably. Comparisons between studies are
290 challenging as there are differences in live weights and slaughter age between countries, leading
291 to differences in feed requirements and dry matter intake. Suckler cows are feed a large
292 proportion grass silage and pasture in both Norway and the other Scandinavian countries
293 (Mogensen et al., 2015). Similar to the semi-intensive production system in Norway, the
294 intensive system in Sweden and Denmark have an intensive finishing of bull calves with approx.
295 50% concentrates, whereas the proportion concentrates in heifer diets have more variation
296 dependent on country and feeding intensity (Mogensen et al., 2015). The Irish and extensive beef
297 production system in Denmark have a larger proportion pasture, and lower proportion of
298 concentrates in the diet compared with average Norwegian beef production (Foley et al., 2011;
299 Mogensen et al., 2015).

300 In flatlands for both breeds and mountains for the continental breeds, C sequestration had
301 a mitigating effect on the emission intensity of beef production. The C mitigation was from the
302 sequestration of manure, feed production and use of pasture. The British breeds produce less
303 manure (due to lower DMI and body weight), which increases the use of synthetic fertilizer and
304 reduces C sequestration. Soussana et al. (2007) concluded that European grasslands are likely to
305 act as atmospheric C sinks, which underlines the importance of including C sequestration in the
306 estimations of emission intensities from pastoral beef production systems.

307 Some whole-farm models, such as Irish BEEFGEM model (Foley et al., 2011), do not
308 include C changes because the C sequestration in soils cannot continue indefinitely. As soil C
309 builds, its decay also increases, and as rate of decay approaches rate of input, soil C reaches an
310 approximate steady state (Guyader et al., 2016). By excluding the soil C change from our
311 estimates, the emission intensities increase to 29.63-31.70 CO₂ eq (kg carcass)⁻¹ for the average

312 farms (Table 8). When excluding soil C change the differences between locations decreased,
313 which indicates that the inclusion of soil C in the calculation of emission intensities can have a
314 marked effect on the outcome, especially for pastoral based beef production systems. The studies
315 of beef production in Denmark and Sweden included the contribution from soil C changes based
316 on the Bern Carbon Cycle Model of Petersen et al. (2013). The Bern Carbon Cycle Model
317 quantifies the change in CO₂ in the atmosphere based on C added to the soil, the release of CO₂
318 from the soil and the decay of C. In Denmark and Sweden the contribution from C sequestration
319 were from -1.8 to -2.4 CO₂ eq (kg carcass)⁻¹ (Mogensen et al., 2015). This is within the range of
320 the level of C sequestration found in the present study of 0.31 to -2.13 CO₂ eq (kg carcass)⁻¹.

321 The Continental breeds are heavier, have a higher feed requirement, and thus produce
322 more enteric CH₄. However, they also have a higher slaughter weight and produce more beef,
323 thus emission intensity is lower. The location will dictate the use of pastures and can influence
324 enteric CH₄ emissions through feed quality and C sequestration through soil, weather and use of
325 inputs. In accordance with White et al. (2010), who reported average GHG emission intensities
326 from beef production systems in New Zealand of 26.0 CO₂ eq (kg carcass)⁻¹ from lowlands and
327 34.0 CO₂ eq (kg carcass)⁻¹ in uplands, our estimates imply that location, farm size, resources and
328 climatic conditions of the farm is important when estimating emission intensities. The locations
329 in the present paper differ in both average temperatures and areas available for crop and silage
330 production, cultivated pastures and outfield pastures (Table 2). The different climatic zones and
331 altitudes influence the production conditions as well as crop and grass yields. By keeping the
332 animal numbers and kg carcass produced constant within breed in the present paper, the emission
333 intensities estimated can be interpreted in the context of location. Flatlands has higher soil N₂O
334 and energy CO₂ emissions than mountains due to greater crop production and use of input factors

335 such as fuel and fertilizer. However, greater crop and grass production in flatlands combined
336 with favorable soil and weather conditions gives greater higher C sequestration compared with
337 mountains. The sensitivity analysis indicate that the emission intensities are dependent on the
338 farm and herd size within location in addition to resources and climatic condition as the emission
339 intensities increase when farm size is reduced.

340 HolosNorBeef does not include aspects of sustainability beyond GHG emissions, which
341 is important to consider in the climate debate. Suckler cow beef accounts for approx. 30% of the
342 beef production in Norway (Animalia, 2018) and the remaining 70% are from dual purpose milk
343 and beef production. The use of pastoral systems have several advantages (i.e., reduced feed
344 costs, animal welfare, carbon sequestration, maintenance of landscape) and grazing preserves
345 biodiversity (Luoto et al., 2003 as cited by Mogensen et al., 2015; Guyader et al., 2016) as well
346 as increases the albedo effect (Kirschbaum et al., 2011). The ecosystems services provided by
347 pastoral beef production systems are not captured by models estimating GHG intensities.

348 The scenarios examined in the present study estimate average emissions based on average
349 farms and management practices, disregarding uncertainties associated with the input data as the
350 use of average farms give a transparent evaluation of the model. Use of average farm scenarios
351 for estimating GHG emissions has limitations, and does not account for the variation in
352 production systems, choice of breed due to resource base, management practices, feeds and feed
353 quality. Future uses of the model will estimate the emission intensities from actual farms
354 distributed geographically across Norway.

355 **5. Conclusions**

356 The whole-farm approach estimated emission intensities of 27.5-32.0 CO₂ eq (kg carcass)⁻¹ from
357 typical herds of British and Continental breeds in two geographically different regions. When
358 excluding soil C the difference between locations decreased in terms of GHG emission intensity,
359 which imply that geographical location is important to consider when estimating emission
360 intensities. Soil C changes must be included in the model for a more a more complete assessment
361 of GHG intensity of beef production from pastoral systems.

362 **Acknowledgements**

363 The authors would like to thank the Norwegian University of Life Sciences and Department of
364 Animal and Aquacultural Sciences for funding the PhD project.

365 **References**

- 366 Alemu, A.W., Amiro, B.D., Bittman, S., MacDonald, D., Ominski, K.H., 2017. Greenhouse gas
367 emission of Canadian cow-calf operations: A whole farm assessment of 295 farms. *Agric.*
368 *Syst.* 151, 73-83.
- 369 Andrén, O., Kätterer, T., 1997. ICBM: The introductory carbon balance model for exploration of
370 soil carbon balances. *Ecol. Appl.* 7, 1226-1236.
- 371 Andrén, O., Kätterer, T., Karlsson, T., 2004. ICBM regional model for estimations of dynamics
372 of agricultural soil carbon pools. *Nutr. Cycl. Agroecosys.* 70, 231-239.
- 373 Animalia, 2017a. Slaughter statistics.
- 374 Animalia, 2017b. Norwegian Beef Cattle Recording System - Annual Report 2016. Norwegian
375 Meat and Poultry Research Centre, Oslo.
- 376 Animalia, 2018. Kjøttets tilstand. Status i norsk kjøtt- og eggproduksjon. Norwegian Meat and
377 Poultry Research Centre, Oslo (In Norwegian).
- 378 Audsley, E., Stecey, K., Parsons, D.J., Williams, A.G., 2009. Estimation of the Greenhouse Gas
379 Emissions from Agricultural Pesticide Manufacture and Use. Granfield University,
380 Bedford, UK. 20pp.
- 381 Beauchemin, K.A., Janzen, H.H., Little, S.M., McAllister, T.A., McGinn, S.M., 2010. Life cycle
382 assessment of greenhouse gas emissions from beef production in western Canada: A case
383 study. *Agric. Syst.* 103, 371-379.
- 384 Beauchemin, K.A., Janzen, H.H., Little, S.M., McAllister, T.A., McGinn, S.M., 2011. Mitigation
385 of greenhouse gas emissions from beef production in western Canada - Evaluation using
386 farm-based life cycle assessment. *Anim. Feed Sci. Technol.* 166-67, 663-677.
- 387 Berglund, M., Cederberg, C., Clason, C., Henriksson, M., Törner, L., 2009. Jordbrukets
388 klimatpåverkan-underlag för att bärekena växthusgasutsläpp på gårdsnivå og

389 nulägesanalyser av exempelgårdar. Delrapport i JOKER-prosjektet.
 390 Hushållningssällskapet, Sweden. 117p (in Swedish).
 391 Bonesmo, H., Skjelvag, A.O., Janzen, H.H., Klakegg, O., Tveito, O.E., 2012. Greenhouse gas
 392 emission intensities and economic efficiency in crop production: A systems analysis of
 393 95 farms. *Agric. Syst.* 110, 142-151.
 394 Bonesmo, H., Beauchemin, K.A., Harstad, O.M., Skjelvag, A.O., 2013. Greenhouse gas
 395 emission intensities of grass silage based dairy and beef production: A systems analysis
 396 of Norwegian farms. *Livest. Sci.* 152, 239-252.
 397 Crosson, P., Shalloo, L., O'Brien, D., Lanigan, G.J., Foley, P.A., Boland, T.M., Kenny, D.A.,
 398 2011. A review of whole farm systems models of greenhouse gas emissions from beef
 399 and dairy cattle production systems. *Anim. Feed. Sci. Technol.* 166-67, 29-45.
 400 Dalgaard, R., Schmidt, J., Halberg, N., Christensen, P., Thrane, M., Pengue, W.A., 2008. LCA of
 401 Soybean Meal. *Int. J. LCA* 13, 240-254.
 402 DNV, 2010. Verification statement no. 76265-2010-OTH-NOR. Yara International ASA. Det
 403 Norske Veritas Certification, Høvik, Norway.
 404 Eurofins, 2015. Feed analysis 2010-2015.
 405 European Commission, 2014. Communication from the Commission to the European Parliament,
 406 the Council, the European Economic and Social Committee and the Committee of the
 407 regions - A policy framework for climate and energy in the period from 2020 to 2030.
 408 European Environment Agency, 2017. Annual European Union greenhouse gas inventory 1990–
 409 2015 and inventory report 2017.
 410 FAO, 2006. World Agriculture: Towards 2030/2050. Food and Agriculture Organization of the
 411 United Nations. Rome.
 412 FAO, 2017. The future of food and agriculture - Trends and challenges. Rome.
 413 Flysjö, A., Cederberg, C., Strid, I., 2008. LCA-databas för konventionella fodermedel [LCA data
 414 regarding conventional animal feed products]. SIK-report 772. The Swedish Institute for
 415 Food and Biotechnology, Göteborg, Sweden. 125 pp (in Swedish).
 416 Foley, P.A., Crosson, P., Lovett, D.K., Boland, T.M., O'Mara, F.P., Kenny, D.A., 2011. Whole-
 417 farm systems modelling of greenhouse gas emissions from pastoral suckler beef cow
 418 production systems. *Agric. Ecosyst. Environ.* 142, 222-230.
 419 Förtabellen, 2008.
 420 Gerber, P.J., Steinfeld, H.H., B. Mottet, A., Opio, C., Dijkman, J., Falcucci, A., Tempio, G.,
 421 2013. Tackling climate change through livestock - A global assessment of emissions and
 422 mitigation opportunities. Food and Agriculture Organization of the United Nations
 423 (FAO), Rome.
 424 Guyader, J., Janzen, H.H., Kroebel, R., Beauchemin, K.A., 2016. Invited Review: Forage
 425 utilization to improve environmental sustainability of ruminant production. *J. Anim. Sci.*
 426 94, 3147-3158. doi: 10.2527/jas.2015-0141
 427 IPCC, 2006. Guidelines for national greenhouse gas inventories. In: Eggleston, H.S., Buendia,
 428 L., Miwa, K., Ngara, T., Tanabe, K. (Eds.), Prepared by the National Greenhouse Gas
 429 Inventories Programme, IGES, Japan. <[http://www.ipcc-nggip-
 430 iges.or.jp/public/2006gl/index.htm](http://www.ipcc-nggip-iges.or.jp/public/2006gl/index.htm)>.
 431 Janzen, H.H., Beauchemin, K.A., Bruinsma, Y., Campbell, C.A., Desjardins, R.L., Ellert, B.H.,
 432 Smith, E.G., 2003. The fate of nitrogen in agroecosystems: An illustration using
 433 Canadian estimates. *Nutr. Cycl. Agroecosyst.* 67, 85-102.

434 Katterer, T., Andersson, L., Andren, O., Persson, J., 2008. Long-term impact of chronosequential
435 land use change on soil carbon stocks on a Swedish farm. *Nutr. Cycl. Agroecosys.* 81,
436 145-155.

437 Katterer, T., Andren, O., 2009. Predicting daily soil temperature profiles in arable soils in cold
438 temperate regions from air temperature and leaf area index. *Acta. Agr. Scand. B-S P.* 59,
439 77-86.

440 Kirschbaum, M. U. F., Whitehead, D., Dean, S. M., Beets, P. N., Shepherd, J. D., and Ausseil,
441 A.-G. E.: Implications of albedo changes following afforestation on the benefits of forests
442 as carbon sinks, *Biogeosciences*, 8, 3687-3696, <https://doi.org/10.5194/bg-8-3687-2011>,
443 2011. Little, S., Lindeman, J., Maclean, K., Janzen, H.H., 2008. HOLOS. A tool to
444 estimate and reduce greenhouse gases from farms. Methodology and algorithms for
445 version 1.1.x. Agriculture and Agri-Food Canada, Cat. No. A52-136/2008E-PDF.

446 Luoto, M., Rekolainen, S., Aakkula, J., Pykala, J., 2003. Loss of plant species richness and
447 habitat connectivity in grasslands associated with agricultural change in Finland. *Ambio.*
448 32, 447-452.

449 Microsoft Corporation, 2016. Available online at <http://microsoft.com/excel>. One Microsoft
450 Way, Redmond, Washington, USA.

451 Mogensen, L., Kristensen, T., Nielsen, N.I., Spleth, P., Henriksson, M., Swensson, C., Hessle,
452 A., Vestergaard, M., 2015. Greenhouse gas emissions from beef production systems in
453 Denmark and Sweden. *Livest. Sci.* 174, 126-143.

454 Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.-F.
455 Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura and
456 H. Zhang, 2013: Anthropogenic and Natural Radiative Forcing. In: *Climate Change*
457 *2013: The Physical Science Basis. Contribution of Working Group I to the Fifth*
458 *Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D.*
459 *Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and*
460 *P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and*
461 *New York, NY, USA.*

462 Nguyen, T.L.T., Hermansen, J.E., Mogensen, L., 2010. Environmental consequences of different
463 beef production systems in the EU. *J. Clean. Prod.* 18, 756-766.

464 NIBIO, 2015. Account Results in Agriculture and Forestry 2015. Norsk institutt for bioøkonomi,
465 Ås, Norway. 247 pp.

466 NIBIO, 2016. Totalkalkylen for jordbruket. Jordbrukets totalregnskap 2014 og 2015. Budsjett
467 2016, in: *Budsjettnemda for jordbruket (Ed.)*.(in Norwegian)

468 NIBIO, 2018. Kilden.
469 <<https://kilden.nibio.no/?X=7334000.00&Y=400000.00&zoom=0&lang=nb&topic=areal>
470 [informatjon&bgLayer=graatone_cache](https://kilden.nibio.no/?X=7334000.00&Y=400000.00&zoom=0&lang=nb&topic=areal)>.

471 Norwegian Meteorological Insitute and Det norske hageselskap, 2006. Klimasonkart (In
472 Norwegian).

473 NRK and Norwegian Meteorological Insitute, 2018. The climate in Norway and the world.

474 Petersen, B.M., Knudsen, M.T., Hermansen, J.E., Halberg, N., 2013. An approach to include soil
475 carbon changes in life cycle assessments. *J. Clean. Prod.* 52, 217-224.

476 Refsgaard Andersen, H., 1990. Ammekoens energibehov og foderoptagelseskapacitet. National
477 Institute of Animal Science, Denmark.

478 Schils, R.L.M., Olesen, J.E., del Prado, A., Soussana, J.F., 2007. A review of farm level
479 modelling approaches for mitigating greenhouse gas emissions from ruminant livestock
480 systems. *Livest. Sci.* 112, 240-251.

481 Skjelvåg, A.O., Arnoldussen, A.H., Klakegg, O., Tveito, O.E., 2012. Farm specific natural
482 resource base data for estimating greenhouse gas emissions. *Acta Agr. Scand. A: Anim.*
483 *Sci.* 62, 310-317.

484 Soussana, J.F., Allard, V., Pilegaard, K., Ambus, P., Amman, C., Campbell, C., Ceschia, E.,
485 Clifton-Brown, J., Czobel, S., Domingues, R., Flechard, C., Fuhrer, J., Hensen, A.,
486 Horvath, L., Jones, M., Kasper, G., Martin, C., Nagy, Z., Neftel, A., Raschi, A., Baronti,
487 S., Rees, R.M., Skiba, U., Stefani, P., Manca, G., Sutton, M., Tubaf, Z., Valentini, R.,
488 2007. Full accounting of the greenhouse gas (CO₂, N₂O, CH₄) budget of nine European
489 grassland sites. *Agric. Ecosyst. Environ.* 121, 121-134.

490 Sozanska, M., Skiba, U., Metcalfe, S., 2002. Developing an inventory of N₂O emissions from
491 British soils. *Atmos. Environ.* 36, 987-998.

492 Statistics Norway, 2017. < <https://www.ssb.no/jord-skog-jakt-og-fiskeri/statistikker/korn/aar>>
493 The Norwegian Environment Agency, 2017. Greenhouse Gas Emissions 1990-2015, National
494 Inventory Report.

495 White, T.A., Snow, V.O., King, W.M., 2010. Intensification of New Zealand beef farming
496 systems. *Agric. Syst.* 103, 21-35.

497 Zhang, W.F., Dou, Z.X., He, P., Ju, X.T., Powlson, D., Chadwick, D., Norse, D., Lu, Y.L.,
498 Zhang, Y., Wu, L., Chen, X.P., Cassman, K.G., Zhang, F.S., 2013. New technologies
499 reduce greenhouse gas emissions from nitrogenous fertilizer in China. *P. Natl. Acad. Sci.*
500 *USA* 110, 8375-8380.

501 Öko-Institut, 2010. E.V., Tankstelle\Diesel-DE-2010.
502 <[http://www.probas.umweltbundesamt.de/php/volltextsuche.php?&prozessid=%7b9F010](http://www.probas.umweltbundesamt.de/php/volltextsuche.php?&prozessid=%7b9F010C0D-A18D-4163-B86D-22E8656276F7%7d&id=1&step=1&search=Tankstelle/Diesel-DE&b=1#in-output)
503 [C0D-A18D-4163-B86D-22E8656276F7%7d&id=1&step=1&search=Tankstelle/Diesel-](http://www.probas.umweltbundesamt.de/php/volltextsuche.php?&prozessid=%7b9F010C0D-A18D-4163-B86D-22E8656276F7%7d&id=1&step=1&search=Tankstelle/Diesel-DE&b=1#in-output)
504 [DE&b=1#in-output](http://www.probas.umweltbundesamt.de/php/volltextsuche.php?&prozessid=%7b9F010C0D-A18D-4163-B86D-22E8656276F7%7d&id=1&step=1&search=Tankstelle/Diesel-DE&b=1#in-output)>. (in German).

505 Åby, B.A., Aass, L., Sehested, E., Vangen, O., 2012. A bio-economic model for calculating
506 economic values of traits for intensive and extensive beef cattle breeds. *Livest. Sci.* 143,
507 259-269.

508 Åby, B.A., Kantanen, L., Aass, L., Meuwissen, T., 2014. Current status of livestock production
509 in the Nordic countries and future challenges with a changing climate and human
510 population growth, *Acta Agr. Scand. A: Anim. Sci.* 64:2, 73-97.

511

Gas/source	Emission factor/equation	Reference
<i>Methane</i>		
Enteric fermentation	$(0.065/55.64) \text{ kg CH}_4 \text{ (MJ GEI)}^{-1}$	(IPCC, 2006)
Relative effect of digestibility (DE%) of feed	$0.1058 - 0.006 \times DE$	(Bonesmo et al., 2013)*
Max.CH ₄ producing capacity of manure (B _o)	$0.18 \text{ m}^3 \text{ CH}_4 \text{ kg}^{-1}$	(IPCC, 2006)
Deep bedding manure	$0.17 \text{ kg CH}_4 \text{ (VS)}^{-1}$	(IPCC, 2006)
Pasture manure	$0.01 \text{ kg CH}_4 \text{ (VS)}^{-1}$	(IPCC, 2006)
<i>Direct nitrous oxide</i>		
Soil N inputs**	$0.01 \text{ kg N}_2\text{O-N (kg N)}^{-1}$	(IPCC, 2006)
Relative effect of soil water filled pore space (WFPS mm)	$0.4573 + 0.01102 \times \text{WFPS}$	(Sozanska et al., 2002)***, (Bonesmo et al., 2012)***
Relative effect of soil temperature at 30cm (ts30°C)	$0.5862 + 0.03130 \times \text{ts30}$	(Sozanska et al., 2002)***, (Bonesmo et al., 2012)***
Deep bedding manure	$0.01 \text{ kg N}_2\text{O-N (kg N)}^{-1}$	(IPCC, 2006)
Pasture manure	$0.02 \text{ kg N}_2\text{O-N (kg N)}^{-1}$	(IPCC, 2006)
<i>Indirect nitrous oxide</i>		
Soil N inputs**	Leaching:	
	EF= $0.0075 \text{ kg N}_2\text{O-N (kg N)}^{-1}$,	(IPCC, 2006),
	Frac _{leach} = $0.3 \text{ kg N (kg N)}^{-1}$	(Little et al.,
	Volatilization:	2008)****
Deep bedding manure	EF= $0.01 \text{ kg N}_2\text{O-N (kg N)}^{-1}$,	
	Frac _{volatilization} = $0.1 \text{ kg N (kg N)}^{-1}$	(IPCC, 2006)
Deep bedding manure	Leaching:	
	EF= $0.0075 \text{ kg N}_2\text{O-N (kg N)}^{-1}$,	(IPCC, 2006)

	$\text{Frac}_{\text{leach}}=0 \text{ kg N (kg N)}^{-1}$	
	Volatilization:	
	$\text{EF}= 0.01 \text{ kg N}_2\text{O-N (kg N)}^{-1}$,	(IPCC, 2006)
	$\text{Frac}_{\text{volatilization}}=0.3 \text{ kg N (kg N)}^{-1}$	
Pasture manure	Leaching:	
	$\text{EF}= 0.0075 \text{ kg N}_2\text{O-N (kg N)}^{-1}$,	(IPCC, 2006),
	$\text{Frac}_{\text{leach}} 0.3 \text{ kg N (kg N)}^{-1}$	(Little et al.,
	Volatilization:	2008)****
	$\text{EF}= 0.01 \text{ kg N}_2\text{O-N (kg N)}^{-1}$,	(IPCC, 2006)
	$\text{Frac}_{\text{volatilization}}=0.2 \text{ kg N (kg N)}^{-1}$	
<i>Soil carbon</i>		
Young (ky) soil C decomposition rate	0.8 year^{-1}	(Andrén et al., 2004)
Old (ko) soil C decomposition rate	0.007 year^{-1}	(Andrén et al., 2004)
Humification coefficient (h) of grass and crop residue	0.13	(Katterer et al., 2008)
Humification coefficient (h) of cattle manure	0.31	(Katterer et al., 2008)
<i>Direct carbon dioxide</i>		
Diesel fuel use	$2.7 \text{ kg CO}_2 \text{ L}^{-1}$	(The Norwegian Environment Agency, 2017)
<i>Indirect carbon dioxide</i>		
Manufacturing N-based synthetic compound fertilizer	$4 \text{ kg CO}_2\text{eq (kg N)}^{-1}$	(DNV, 2010)
Manufacturing pesticides	$0.069 \text{ kg CO}_2\text{eq (MJ pesticide energy)}^{-1}$	(Audsley et al., 2009)
Manufacturing silage additives	$0.72 \text{ kg CO}_2\text{eq (kg CH}_2\text{O}_2)^{-1}$	(Flysjö et al., 2008)

Production of diesel fuel	0.3 kg CO ₂ eq L ⁻¹	(Öko-Institut, 2010)
Production of electricity	0.11 kg CO ₂ eq kWh ⁻¹	(Berglund et al., 2009)
Purchased soya meal	0.93 kg CO ₂ eq (kg DM) ⁻¹	(Dalgaard et al., 2008)
Purchased barley grain	0.62 kg CO ₂ eq (kg DM) ⁻¹	(Bonesmo et al., 2012)

514 GEI= Gross energy intake; VS = volatile solids; WFPS = water filled pore space; ts30 = soil

515 temperature at 30cm; EF = emission factor; Frac_{leach} = Leaching fraction; Frac_{volatilization} =

516 Volatilization fraction

517 *Equation derived by Bonesmo et al. (2013) based on IPCC (2006), Little et al. (2008) and

518 Beauchemin et al. (2010).

519 **Includes land applied manure, grass and crop residue, synthetic N fertilizer, mineralized N

520 ***Equation derived by Bonesmo et al. (2012) using data from Sozanska et al. (2002)

521 ****Value simplified from equation given by Little et al. (2008)

522 *Table 2 Average temperatures (C°) with min and max temperatures (in parenthesis) and land*
 523 *resources (ha) with proportion of total area (in parentheses) from two different locations*
 524 *(flatlands and mountains) in Norway.*

	Flatlands	Mountains
Climatic zone ^a	4*	7*
<i>Average temperatures</i>		
Spring (C°) ^a	6.2 (-13.6;30.7)	5.3 (-15;20.7)
Summer (C°) ^a	14.4 (1.9;25.0)	11.1 (0.1;24.5)
Fall (C°) ^a	5.6 (-9.4;18.6)	4.1 (-17.6;18.4)
Winter (C°) ^a	-5.6 (-25.2;8.9)	-4.2 (-22;10.1)
<i>Land resources</i>		
Cultivated land/cropland (ha) ^b	16,466 (0.13 ^{**})	4,273 (0.02 ^{**})
Cultivated pastures (ha) ^b	3,288 (0.02 ^{**})	3,964 (0.02 ^{**})
Forest (ha) ^b	70,333 (0.55 ^{**})	36,627 (0.16 ^{**})
Bare land (ha) ^b	7,335 (0.06 ^{**})	161,558 (0.71 ^{**})
Rich vegetation (ha) ^b	3,223 (0.44 ^{***})	40,258 (0.25 ^{***})
Medium rich vegetation (ha) ^b	734 (0.10 ^{***})	39,369 (0.24 ^{***})
Poor vegetation (ha) ^b	41 (0.01 ^{***})	52,842 (0.33 ^{***})
Bare mountain (ha) ^b	0 (0.00 ^{***})	20,688 (0.13 ^{***})
Unclassified (ha) ^b	3,337 (0.45 ^{***})	8,400 (0.05 ^{***})

525 ^a NRK and Norwegian Meteorological Insitute (2018)

526 ^b Norwegian Institute of Bioeconomy Research (NIBIO, 2018)

527 * On a scale from 1 (good) to 4 (harsh)

528 ** Do not sum up to 100% as area unrelated to agriculture are left out of the table

529 *** Proportion of bare land.

530 *Table 3 Average animal data for Norwegian beef farms used to estimate GHG emission*
 531 *intensities in two locations.*

Farm characteristics (unit)	British	Continental
Beef produced (kg carcass) ^{ab}	7699	9635
Cows, average weight (kg LW) ^c	600	800
Cows, carcass weight (kg) ^c	324	432
Cows, concentrate (proportion) ^c	0.25	0.17
Cows, time on pasture (proportion) ^c	0.36*	0.38**
Milk, yield (kg raw milk year ⁻¹) ^c	1,100	1,600
Twinning frequency (%) ^a	1.9	3.0
Still born (%) ^a	3.5	3.9
Dead before 180 days (%) ^a	3.6	4.1
Gender distribution (proportion heifers) ^c	0.5	0.5
Heifers, birth weight (kg LW) ^c	38	42
Heifers, weaning weight (kg LW) ^c	251	295
Heifers, yearling weight (kg LW) ^c	365	416
Heifers, carcass weight (kg) ^c	206	244
Heifers, age at slaughter (month) ^a	18.2	17.5
Heifers, age at first calving (month) ^c	26.5	28.9
Heifers, concentrate birth-slaughter (proportion) ^c	0.22	0.38
Heifers, time on pasture (proportion) ^c	0.19	0.13
Young bulls, birth weight (kg LW) ^c	40	45
Young bulls, weaning weight (kg LW) ^c	269	322
Young bulls, yearling weight (kg LW) ^c	445	547
Young bulls, carcass weight (kg) ^a	291	353
Young bulls, age at slaughter (month) ^a	17.5	16.8
Young bulls, concentrate birth-slaughter (proportion) ^c	0.53	0.50

532 LW= live weight

533 ^a Animalia (2017a)

534 ^b Norwegian Institute of Bioeconomy Research (NIBIO, 2015)

535 ^c Åby et al. (2012)

536 *42% cultivated pasture, 58% outfield pasture

537 **50% cultivated pasture, 50% outfield pasture

538 *Table 4 Average animal numbers, crop and fuel usage data for Norwegian beef farms used to*
 539 *estimate GHG emission intensities from two different locations (flatlands and mountains) in*
 540 *Norway.*

Farm characteristics	Flatlands	Mountains
<i>Animal system</i>		
Cows (year ⁻¹) ^a	28	28
Calves born (year ⁻¹) ^a	28	28
Replacement heifers (year ⁻¹) ^a	10	10
Heifers slaughtered (year ⁻¹) ^a	4	4
Young bulls slaughtered (year ⁻¹) ^a	13	13
<i>Input use</i>		
Fuel (L year ⁻¹) ^a	3854	2947
Electricity (kWh year ⁻¹) ^a	26300	29100
Silage additive (kg CH ₂ O ₂ year ⁻¹) ^b	803	416
Ley synthetic fertilizer (kg N ha ⁻¹) ^b	13	13
Ley pesticide (MJ ha ⁻¹) ^a	1.1	1.1
Barley synthetic fertilizer (kg N ha ⁻¹) ^b	9.5	9.5
Barley pesticide (MJ ha ⁻¹) ^a	29.8	29.1
Oats synthetic fertilizer (kg N ha ⁻¹) ^b	8.5	8.5
Oats pesticide (MJ ha ⁻¹) ^a	14.5	14.1
Spring wheat synthetic fertilizer (kg N ha ⁻¹) ^b	10	10
Spring wheat pesticide (MJ ha ⁻¹) ^a	34.1	33.2
Winter wheat synthetic fertilizer (kg N ha ⁻¹) ^b	12.1	12.1
Winter wheat pesticide (MJ ha ⁻¹) ^a	64.1	64.1
<i>Land use</i>		
Farm size (ha) ^a	44.6	41.5
Pasture and ley area (ha) ^a	38.9	40.1
Grass yield (FUm/ha) ^a	3020	3190
Grass silage nutritive value (FUm) ^c	0.87	0.84
Barley area (ha) ^{ad}	3.0	0.9

Barley yield (kg DM ha ⁻¹) ^{ade}	4310	2840
Oats area (ha) ^{ad}	1.5	0.1
Oats yield (kg DM ha ⁻¹) ^{ade}	4030	2960
Spring wheat area (ha) ^{ad}	1.1	0.0
Spring wheat yield (kg DM ha ⁻¹) ^{ade}	4860	3870
Winter wheat area (ha) ^{ad}	0.1	0.0
Winter wheat yield (kg DM ha ⁻¹) ^{ade}	4860	3870

541 FUm = feed units milk

542 ^a Norwegian Institute of Bioeconomy Research (NIBIO, 2015)

543 ^b Norwegian Institute of Bioeconomy Research (NIBIO, 2016)

544 ^c Eurofins (2015)

545 ^d Statistics Norway (2017)

546 ^e Fôrtabellen (2008)

547 *Table 5 Natural resource data used to estimate GHG emission intensities from two different*
 548 *locations (flatlands and mountains) in Norway (Bonesmo et al., 2013; Skjelvåg et al., 2012).*

	Flatlands		Mountains	
	Grassland	Field crops	Grassland	Field crops
Soil temperature at 30 cm depth, winter (°C) ^a	-0.68	-0.67	-0.39	0.90
Soil temperature at 30 cm depth, spring (°C) ^a	5.37	5.16	3.85	6.67
Soil temperature at 30 cm depth, summer (°C) ^a	13.79	13.80	10.81	13.93
Soil temperature at 30 cm depth, fall (°C) ^a	5.20	5.16	4.05	6.95
Water filled pore space, winter (%) ^b	65	65	74	68
Water filled pore space, spring (%) ^b	48	51	57	55
Water filled pore space, summer (%) ^b	43	48	45	51
Water filled pore space, fall (%) ^b	62	65	65	68
$r_w \times r_T$ yearly (dimensionless) ^c	0.94	1.06	0.65	1.29
Soil organic C (Mg ha ⁻¹)	6		8	

549 ^a Estimated according to Katterer and Andren (2009).

550 ^b Estimated according to Bonesmo et al. (2012).

551 ^c Estimated according to Andrén et al. (2004).

552 *Table 6 Average animal numbers, carcass production, land use and farm inputs for small and*
 553 *large farms of British breeds located in the flatlands used to investigate the sensitivity to*
 554 *variation in farm size and corresponding impact on GHG emission intensities compared with the*
 555 *average farm*.*

Farm characteristics	Small farm	Large farm
<i>Animal system</i>		
Cows (year ⁻¹) ^a	14.4	38
Calves born (year ⁻¹) ^a	14.4	40
Replacement heifers (year ⁻¹) ^a	5	14
Heifers slaughtered (year ⁻¹) ^a	2	5
Young bulls slaughtered (year ⁻¹) ^a	7	19
Beef produced (kg carcass) ^{ab}	3946	10851
<i>Input use</i>		
Fuel (L year ⁻¹) ^a	2071	5729
Electricity (kWh year ⁻¹) ^a	18300	38200
Silage additive (kg CH ₂ O ₂ year ⁻¹) ^c	323	593
<i>Land use</i>		
Farm size (ha) ^a	25.1	74.8
Pasture and ley area (ha) ^a	24.6	63.3
Barley area (ha) ^{ad}	0.2	5.9
Oats area (ha) ^{ad}	0.1	3.0
Spring wheat area (ha) ^{ad}	0.1	2.1
Winter wheat area (ha) ^{ad}	0.0	0.9

556 **Factors not included are similar to the baseline, British breeds located in the flatland*

557 ^a Norwegian Institute of Bioeconomy Research (NIBIO, 2015)

558 ^b Animalia (2017a)

559 ^c Norwegian Institute of Bioeconomy Research (NIBIO, 2016)

560 ^d Statistics Norway (2017)

561 *Table 7 Emissions and proportion of total emissions (in parenthesis) from average herds of*
 562 *British and Continental breeds in two different locations (flatlands and mountains) in Norway*
 563 *(kg CO₂ eq).*

	Flatlands		Mountains	
	British	Continental	British	Continental
Enteric CH ₄	108,011 (0.47)	127,729 (0.48)	108,307 (0.44)	128,091 (0.45)
Manure CH ₄	24,814 (0.11)	30,532 (0.12)	25,054 (0.10)	30,823 (0.11)
Manure N ₂ O	23,176 (0.10)	26,835 (0.10)	23,384 (0.9)	27,068 (0.09)
Soil N ₂ O	25,145 (0.11)	29,059 (0.11)	23,713 (0.10)	27,108 (0.10)
Soil C	-13,574 (-0.06)	-20,524 (-0.08)	2,381 (0.01)	-3,046 (-0.01)
Off-farm barley	6,526 (0.03)	11,895 (0.04)	12,638 (0.05)	18,266 (0.06)
Off-farm soya	10,658 (0.05)	16,772 (0.06)	14,516 (0.06)	20,229 (0.07)
Indirect energy	25,065 (0.11)	25,065 (0.09)	22,959 (0.09)	22,959 (0.08)
Direct energy	17,645 (0.08)	17,645 (0.07)	13,492 (0.05)	13,492 (0.05)
Total emissions	227,466	265,006	246,445	284,991
Total emissions ex. soil C	241,040	285,531	244,064	288,037

564

565 *Table 8 GHG emission intensities from average herds of British and Continental breeds in two*
 566 *different locations (flatlands and mountains) in Norway (CO₂ eq kg⁻¹ carcass).*

	Flatlands		Mountains	
	British	Continental	British	Continental
Enteric CH ₄	14.03	13.26	14.07	13.29
Manure CH ₄	3.22	3.17	3.25	3.20
Manure N ₂ O	3.01	2.79	3.04	2.81
Soil N ₂ O	3.27	3.02	3.08	2.81
Soil C	-1.76	-2.13	0.31	-0.32
Off-farm barley	0.85	1.23	1.64	1.90
Off-farm soya	1.38	1.74	1.89	2.10
Indirect energy	3.26	2.60	2.98	2.38
Direct energy	2.29	1.83	1.75	1.40
Total emissions	29.54	27.50	32.01	29.58
Total emissions ex. soil C	31.31	29.63	31.70	29.89

567

568 *Table 9 Sensitivity elasticities for the effect of 1% change in the selected emission factors (EF)*
 569 *and initial soil organic carbon on the greenhouse gas (GHG) emission intensities CO₂ eq (kg*
 570 *carcass)⁻¹.*


	Response	% change in CO ₂ eq (kg carcass) ⁻¹
Enteric CH ₄ conversion factor, Y _m	linear	0.47
Manure N ₂ O EF	linear	0.10
IPCC soil N ₂ O EF	linear	0.09
Soil C change external factor ^a	non-linear	0.16
Manufacturing fertilizer EF	linear	0.10
Fuel combined EF	linear	0.09
Initial soil organic carbon	linear	0.12

571 ^a Mean sensitivity elasticity (%) for the change +/- 1% of $r_w \times r_f$.

572 *Table 10 The effect of farm and herd size on the greenhouse gas (GHG) emission intensities CO₂*
 573 *eq (kg carcass)⁻¹.*

	Small farm	Large farm
Enteric CH ₄	14.52	13.50
Manure CH ₄	3.31	3.12
Manure N ₂ O	3.14	2.88
Soil N ₂ O	3.34	3.31
Soil C	-1.49	-1.19
Off-farm barley	1.79	0.43
Off-farm soya	1.92	1.10
Indirect energy	3.63	3.75
Direct energy	2.40	2.42
Total emissions	32.57	29.31
Total emissions (% change from baseline*)	10.12	0.88

574 * Baseline: average herd of British breeds located in the flatlands



575

576 *Figure 1 The suckler cow beef production system.*